
Effect of Red Dragon Fruit Juice on Acrylic Resin Color





(Thank you for your kind assistance & assurance. Kindly revert back within Iweek)

Title: "Effect of Red Dragon Fruit Juice on Acrylic Resin Color"

Article Type: Research Article

Date Sent to Reviewer: 8.3. 2019

Expected Date of Review Completion:

Editor Assigned: Prof. Konstantina Voniati

Editorial Assistant: Dr. Andreas Demopoulos

REVIEW QUESTIONS

1. Title	
Does the title address the topic specified in the paper?	Yes 🗆 No 🗆
Do you recommend unother Title?	CCDINT
34	SCRIPT
2. Abstract(fary) Scien	tific Services
Is the abstract specific and representative of the paper?	Yes □ No □
Are main results and conclusion presented?	Yes □ No □
3. Introduction(if any)	
Is the problem being addressed clearly emphasized?	Yes □ No □
Are the research question and objectives clearly identified?	Yes 🗆 No 🗆
4. Theoretical Foundation	
Do the arguments build on existing theory?	Yes 🗆 No 🗀
Has the author(s) cited the relevant literature?	Yes □ No □
Are propositions/hypotheses important for the area of research?	Yes 🗆 No 🗀
5. Research Design and Methods	
Is the research design adequate?	Yes 🗆 No 🗆
Is the sample size adequate?	Yes □ No □



Is there sufficient detail to enable the reader to duplicate the analysis?	Yes 🗆 No 🗆
6. Results	
Is the statistical analysis clearly presented?	Yes 🗆 No 🗀
Are the results presented in a logical sequence to support the hypotheses?	Yes 🗆 No 🗆
7. Discussion	
Are data and results presented in a succinct manner?	Yes 🗆 No 🗆
Is the relationship between results with previous research relevant?	Yes 🗆 No 🗀
Are reasons for differences in results with previous research clearly stated?	Yes 🗆 No 🗆
Are possible directions for future research stated?	Yes 🗆 No 🗆
8. Conclusions	
Are conclusions stated briefly in a logical order?	Yes 🗆 No 🗅
9. General Aspects	
Is the paper well structured?	Yes □ No □
Is the paper well written and readable?	Yes 🗆 No 🗆
Are references accurate and complete?	Yes 🗆 No 🗆
Is line of reasoning convincing?	Yes 🗆 No 🗆
Would you recommend this paper to your colleagues?	Yes □ No □

GENERAL/SPECIFIC COMMENTS TO THE AUTHOR

For paper improvement please provide in the following constructive comments in the form of a detailed review on the most important aspects to be addressed by the author(s). This part is intended to be communicated to the author(s):

Research Metodology

The type of research used is experimental laboratories with pre and post test with control group design, namely taking measurements after the intervention in the treatment group and comparing the results of measurements with the control group. ----- Please better describe.



Result

Statistical analysis is not clearly presented. The results are not presented in a logical sequence to support the hypotheses. The relationship between the results of previous research is not entirely relevant. The reasons for the differences in results compared to previous research are not clearly stated.

Discussion

A comparison of the findings of the study with other results is completely absent in the Discussion.

Definitions of individual reactions and procedures are included in the discussion, for example:

-----Red dragon fruit juice contains many anthocyanin substances that can give color from colorless to purple.

Conclusion:

The conclusion contains an evaluation of the experiment:

-----An experiment was performed: There were significant differences in the color of the heatcured acrylic resin plate after 7 and 14 days of immersion in distilled water solution and red dragon fruit juice solution. Conclusion: Red dragon fruit juice can cause color discoloration on heat cured acrylic resin based plate.

EVALUATION RESULTS (choose any one)

- 1. Not developed enough and must be "Rejected"
- 2. Revise and resubmit with "Minor Corrections"
- 3. Conditional acceptance with proposed revisions must be completed with "Major Revisions"
- 4. Acceptable in its current form/"Accepted without changes"
- 5. "Strongly Recommended"



Effect of Red Dragon Fruit Juice on Acrylic Resin Color

Fransiska Nuning Kusmawati **, Diah Puspitasari Kusumaningrum b

* Department of Prosthodonsia, Faculty of Dentistry, Universitas Prof DR. Moestopo (Beragama), Jakaria, Indonesia.

^b Clinical Student Department of Prosthodonsia, Faculty of Dentistry, Universitas Prof. DR. Moestopo (Beragama), Jakaria, Indonesia.

Abstract

Background: Discoloration of denture acrylic resin base is one of the problems in appearance for patients who use them. The most common habit can cause discoloration in the denture usually associated with the patients diet. One example is consuming red dragon fruit juice. Red dragon fruit juice contains many anthocyanin substances that can give color from colorless to purple. Furthermore, color changes indenture can also cause by the characteristic of the acrylic resin plate itself which can absorb water due to perosity. Objective: The aim of this study is to investigate the negative effects of consuming dragon fruit juice solution on heat cured acrylic resin based plate. Methods: This type of research is laboratory experimental and the design of this research is pre-test post-test with control group design. The number of samples were sylected using Simple Random Sampling method. The samples were divided into two groups, control (n = 16, distilled water solution) and tentative (n = 16, red dragon fruit juice solution). Color measurement using a digital spectrophotometer (VTTA Eusyshade). The initial color of the plate was measured before immersion into dragon fruit juice and was Prischman statistical test and Mann Whitney statistical test. Findings: There were significant differences in the color of the heat-cured acrylic resin plate after 7 and 14 days of immersion in distilled water solution. Novelv: Red dragon fruit juice can cause color discoloration on heat cured acrylic resin based plate.

Keywords: Red Dragon Fruit Juice Solution; Acrylic Resin Plate Discoloration; Heat-cured Acrylic Resin Base.

1. Introduction

Nowadays, the level of public awareness of dental health is increasing. This allows increased of dentures uses with bases made of acrylic resin [1]. According to the type of polymerization, there are two types of acrylic resins which are heat-cured type and the cold-cured type. Discoloration of acrylic resin can be caused by several factors, one of them is the ability of acrylic resin to absorb liquid. Discoloration of acrylic resin is not only related to physical and chemical properties, but also related to the patient's diet [2].

Hylocereus polyrhizus or red dragon fruit is often called red pitaya because both skin and flesh colors are red. The complete nutritional and vitamin content and phytochemicals in the form of flavonoids in dragon fruit are also known to reduce the risk of cancer. However, the presence of some of these substances might affect the discoloration of the aerylic denture base [3].

1.1. Heat-cured Acrylic Resin

Aerylic resin as a denture base is still being used due to its several advantages including good aesthetic, non-toxic, easy to be manipulated, affordable and easy to be repaired. Among other aerylic resin materials, Polymethyl methaerylate (PMMA) is the most popular aerylic resin material that is used for denture base material [4, 5].

Heat-cured acrylic resins commonly used in dentistry as artificial teeth base. Heat-cured acrylic resin is hot polymerization scrylic resin with heat-activated ingredients. Thermal energy needed for the polymerization of these materials can be obtained by using water heating [6].

One of the drawbacks of using acrylic resin as a denture base is that fractures often occur due to usage, poor thermal conductor, porosity, absorbing liquid, and abration during cleaning. Liquid absorption of heat-cured acrylic resin is also one of the disadvantages of using acrylic resin as a denture base.

There are four types of dragon fruit that have been cultivated in Indonesia, namely white flesh dragon fruit (Hybioceness undatus), pad meat dragon fruit (Hybioceness polyrhizus), super red meat dragon fruit (Hybioceness costariceness), and white flesh syellow dragon fruit (Seleniceness megalanthus). A bundred grams of red dragon fruit (Hybioceness polyrhizus) flesh contains water (82.5-83.0 g); grotein (0.16-0.23 g); fat (0.21-0.01 g); niario (1.20-1.00 g); vitamin (5.00-90 mg); votamin (5.00-90 mg); votamin (5.00-90 mg); votamin (7.70).



Figure 1. Red Dragon Fruit

According to research conducted by Widianingsih, the red dragon fruit also has high autioxidant activity which is 67.45 ppm [11].

67.45 pm; [11].

Anthocyanin is a plant pigment that could be found in nature. This pigment has relevant role in plant propagation, ecophysiology, and plant defines mechanisms and gives color to fruits (red dragon fruit, blueberries, grapes, 6c) and vogetables (spinsel), purple subbage and rosein flowers). Anthocyanin are phenolic compounds containing components that are soluble in water, found in various types of plants and gives color from colorless to purple. Anthocyanin is a class of flavonoid compounds which is widely divided into plant polyphenols and has a group of red to blue pigments that are scattered in plants. The pigments that found in plants have variety of benefits, ²⁸ Blue, rod, and purple pigments extracted from flowers, fruits and vegetables are traditionally used as natural food coloring. Besides being used as natural food coloring agent, some flowers such as rosella flower and red drugon fruit are rich in anthocyanin and have been traditionally used as medicines to treat various diseases [12].

1.2. Color Change Measuring Instrument

The measurement of color changes on heat-cured acrylic resin plate in this study was using digital apectrophotometer (VITA Hasyshade). (Figure 2). This instrument is the latest spectrophotometer used in clinical use



Figure 2. VITA easyshade [14]

Color measurements are based on the use of 3 or 4 votor discs, such or which has been accumulate an important of the color intensity that distinguishes strong colors and weak colors, described the distance of a circle from the center (discussed color ball). What is the color intensity that distinguishes strong colors and weak colors, described as the distance of a circle from the center (discussed color ball). Walse is the color quality associated with lighting, which is the level of brightness and described as a vertical line. The color measurement of this system is to visually match the color of the product with the color of themsel using the cense of sight [13].

2. Research Methodology

The design used was experimental laboratory research with a protest-post test with control group design approach. The endy was conducted in Fessuly of Densitiny, Universities Perol DEL Mosterboy (Beragiams) on July 2018. They two samples of heat-cured acrytic resis plates were prepared with 20 mm length, 10 mm width and 1 mm thickness. Samples were electry by sample random samples.

Samples were estery by simple motion assupting.

The research procedures are earried out as follows: 1) Preparing the tools and materials to be used in the study 2)
Thirty two samples were divided into group 1 and 2 with n number was 16, 3) Performed measurements with digital
peterophotometer (VITA Especial points of the peterophotometer (VITA Especial points) properties from each group (perests), 4) Preparation of pour red dragon fruit
juice solution by blending 168 gr of red dragon fruit with 100 ml of distilled water using juice blender. Juice was thon
poured into the container that has been prepared. 5) Group 1 was immersed into equides, 65 Group 2 was immersed
into red dragon fruit juice. 7) After 7 desys, samples were trinsed with votor, dired and volor checked with digital
9). After 16d of 16 days, camples were removed against sing two access, rinsed with water, dried and color checked with
digital spectrophotometer. 10) Record and compared the measurement results.

3. Resul

5. SCENIII
The statistical text used in this study is the Non Parametric text which resulted in ordinal scale data. The results of
the study were conducted with a descriptive text to study which group had the greatest potential in causing acrylic
resin plate discoloration (Figure 3)



Figure 3. After 14 days immersion

Figure 4 shows the results of chrome samples before and after immersion for 7 and 14 days. These results showed not dropp plate has the generate presental for discoloration on beat-cardon deright result palse. Figure 4 shows the results of value in samples before and after immersion for 7 and 14 days. These results showed red dragon juice has the greatest potential for cooler change in the them-teroid excity it result palse.




Figure 4. Chrome after immersion




Figure 5, Value after immersion

Table 1. Friedman test Chrome

Value	Mean Ranks
Aquades before immersion	1.60
Aquades after immersion 7days	2,34
Aquades after immerson 14 days	2.66
Red dragon juice before immersion	1.00
Red dragon juice after immersion 7 days	2.00
Red dragon juice after immersion 14 daysi	3.00
Aquades	

N	16
Chi-Square	26.881
df	2
Asymp. Sig.	.000
Larutan jus buah naga merah	
N	16
Chi-Square	32.000
df	2
Asymp. Sig.	.000
Value	Mean Ranks
Aquades before immersion	3.00
Aquades after immersion 7 days	1.94
Aquades after immersion 14 days	1.06
Red dragon juice before immersion	3.00
Red dragon juice after immersion 7 days	2.00
Red dragon juice after immersion 14 daysi	1.00
Aquades	
N	16
Chi-Square	30.125
df	2
Asymp, Sig.	.000
Red dragon juice	
N	16
Chi-Square	32,000
df	2
Asymp. Sig.	.000

Friedman's statistical test results for both groups of heat-cured acrylic resin plates in chrome and value colors. Statistical test results on chrome color for distilled water before immersion, 7 days immersion and 14 days immersion showed a value of p=0.000 (p=0.05), meaning that there were significant differences in the color of chrome on heat-cured acrylic resin tools in the control group. Statistical test results on the chrome color for dargon fruit juice solution before immersion, 7 days immersion and 14 days immersion showed a value of p=0.000 (p<0.05), meaning that there were significant differences in the color of chrome in heat-cured acrylic resin tools in the treatment group. Statistical test results on the color value for distilled water before immersion, 7 days immersion and 14 days immersion showed a value of p=0.000 (p<0.05), meaning that there were significant differences in color values on heat-cured acrylic resin tools in the control group. Statistical test results on the color roule of dragon fruit juice solution before immersion, 7 days immersion and 14 days immersion showed a value of p=0.000 (p<0.05), meaning that there were significant differences in color values in the heat-cured acrylic resin tools in the treatment group.

Table 2. Mann Whitney test

Before immersion	
Aquades and	Red dragon juice
Mann-Whitney U	124.500
Wilcoson W	260.500
z	132
Asymp, Sig. (2-tailed)	.895
After imn	erston 7 days
Aquades and	Red dragon juice
Mann-Whitney U	15.500

Wilcoxan W	151.500	
z	-1.266	
Asymp. Sig. (2-tailed)	.000	
After imm	ersion 14 days	
Aquades and	Red dragon juice	
Mann-Whitney U	15.500	
Wilepson W	151.500	
Z	4.266	
Asymp. Sig. (2-tailed)	.000	
Val	ue test	
Before	immersion	
Aquades and	Red dragon juice	
Mann-Whitney U	104,500	
Wilcoxen W	240,500	
Z	-,888	
Asymp. Sig. (2-tailed)	.375	
After imm	ersion 7 days	
Aquades and	Red dragon juice	
Menn-Whitney U	47,000	
Wilcoxon W	183,000	
z	-3.059	
Asymp. Sig. (2-tailed)	.002	
After imm	ersion 14 days	
Aquades and	Red dragon juice	
Mann-Whitney U	0.000	
Wilcoxon W	136.000	
2	-4.828	
Asymp. Sig. (2-tailed)	.000	

The Mann Whitney statistical test results for both groups of heat-cured acrylic resin plates on chrome color and color value. Statistical test results on chrome color for aquades and red dragon fruit juice solution before immersion showed a value of p = 0.995 (p = 0.05), this means that there is so significant difference in the color of the chrome in the heat-cured acrylic each better the immersion of the control and treatment groups. For chrome color in aquades and 7 days immersion red dragon fruit juice solution showed a value of p = 0.197 (p = 0.05), meaning that there were no significant differences in the color of chrome in heat-cured acrylic resit notes in the 7 day immersion of the control and treatment groups. For the chrome color of distilled water and 14 days immersion red dragon fruit juice solution showed a value of p = 0.000 (p = 0.000), meaning that there were significant differences in the color of chrome in heat-cured acrylic resin in the 14 day immersion of the control group and the treatment group. So it can be concluded that there are significant differences in chrome color on the heat-cured acrylic resin plate between groups in the post-lest data (after 14 days immersion).

In this study, each of the heut-cured acrylic resin plates were immersed according to their groups for 7 days and continued for up to 14 days. Determination of immersion time for 7 days refers to research conducted by Turkan, because it is assumed to be identical with the use of artificial teeth from beat-cured acrylic resin by drinkers of red dragon fruit juice for 2 years. Red dragon fruit juice drinker is a person who has a habit of consuming red dragon fruit juice fruits once a day. The estimation time of a person to drink dragon fruit juice is about 15 minutes. Immersion for 7 days is equivalent to 2 years of use, and immersion for 14 days is equivalent to 4 years of use that calculated in equations below.

$$\frac{7 \ days \times 24 \ hours \times 60 \ minutes}{15 \ minutes} = 672 \ days \ (2 \ years)$$

$$\frac{14 \, days \times 24 \, hours \times 60 \, minutes}{15 \, minutes} = 1344 \, days \, (4 \, years)$$

This study uses a heat-cured acrylic resin plate due to the increasing level of public awareness of dental health. This allows increased use of dentures with the basic ingredients made of acrylic resin [1]. Acrylic resin materials have advantages such as montoxic, does not irritate tissue, meet seathetic requirements, relatively cheap prices, easy to be manipulated and repaired. In addition to its beneficial properties, acrylic resins have several disadvantages including porosity which results in absorbing water or liquid, food scraps or chemicals and less abcasion resistance.¹³ Acrylic resin materials have the property of absorbing water gradualty over a period of time with absorption mechanism through the diffusion of water molecules according to the law of diffusion [15].

The results of this study indicated that solution of pure red dragon fruit juice caused discolouration on heat-cured acrylic resin plates. This occurs due to the imbibition process experienced by acrylic resin plates against anthocyanin substances that present in red dragon fruit, This process occurs because polymethyl metallarylate which is the basic material of acrylic resin has a tendency to absorb water through the imbibition process. Since non-crystalline structure has high internal energy, molecular diffusion could occur in the resin.

man angon meneral energy, moscular arraison could occur in the resin.

Color changes of hest-send acyclic resin plate due to immersion in pure red drugon finit juice solution can be caused by several factors. The first factor is the characteristic of the acrylic resin itself which can absorb liquid. The second factor is the intensity of contaming red dragon fruit. The third factor is the promity of acrylic resin which can absorb water or liquid, food contaming red dragon fruit. The third factor is the promity of acrylic resin which can absorb water or liquid, food carrops or chemicals and is less abrasion resistant. The fourth factor is the surface roughness of acrylic resin, if the surface roughness is large it can be a place of accumulation of dye stains that cause color changes.

5. Conclusion and Recommendation

Based on the results of research on the changes in the color of the heat-cured acrylic resin plate after immersion in red dragon fruit juice solution, it can be concluded that color changes that occur on the heat-cured acrylic resin plate can be caused by solution of red dragon fruit juice (Hylocereus polyrhizus). What can be done to progress this research going forward is:

- . Conduct research with a larger sample and experiment with a longer soaking time to get a more accurate final
- · Add a dose to the solution of the red dragon fruit juice with different consistency in order to get more accurate

6. Declaration of Competing Interest


The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

All procedures performed in studies involving human participants were in accordance with the othical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

8. References

- Deperiemen Kesehaten Republik Indonesia, Laporan Riset Kesehatan Dasar Nasional 2013. Jakarta: Badan Penelirian dan Pengerabangan Kesehatan, 149.
- [2] Anusavice KJ. (2012). Science of Dental Materials, 12th Ed. Philadelphia: W.B. Saunders Co.; 237-51.
- [3] Herdini L. (2015). Uji Efektivitas Ekstrak Daging Buah Naga Merah (Hylocoreus polyrhizus) Terhadap Pertumbuhan Candida albicasa Pada Pata Dasar Gigi Tiraan Rosin Akrilik [akripsi]. Yogyakarta: Universitas Gajah Mada. Avullable online: http://ed.repository.ugm.as.id/rdcs.php/mod-penelitian_detail&astb=PenelitianDetail&act-view&typ=html&bulex_id=84395 &cobyek_id=4 (accessed on 12 July 2018).
- [4] Siswanto A. (2007). The Influence of Sooking Acrylic Plate in Leraes Condensation to Strength Transversa, Journal of Indonesian Dent Association 57(3), 97-100.
- [5] Aaonim, Sifat Resin Akrilik (Biologis, Kimia, Fisika), November 2011, Available https://www.scribd.com/document/190182630/Sifat-Resin-Akrilik-Biologis-Kimia-Fisika (accessed on 12 April 2018).

- [6] Craig RG, Powers JM. (2006). Restorative Dental Materials. 12th Ed. St. Louis CV Mosby Company, 513-53.
- [7] Sitorus Zu, Eddy Dahar (2012). Perbaikan Sifat Fisis dan Mekanis Resin Akrilik Polimerisasi Panas Dengan Penambahan Serat Kaca. Dentika Dental Journal. 17 (1), 24-9.
- [8] Haryanto, A. G., Margo, A., Burhan, L. K., Suryatenggara, F., & Setiabudi, I. (2012). Buku ajar gigi geligi tiruan sebagian lepasan jilid 1. Jakarta: Hipokrates, 4-15.
- [9] Handayani, P. A., & Rahmuwati, A. (2012). Pemanfaatan kulit buah naga (dragon fruit) sebagai pewarna alami makanan pengganti pewarna sintetis. Jurnal bahan alam terbarukan, 1(2).
- [10] Kres Dahana, S. P., & Warisno, S. (2013). Buku pintar bertanam buah naga. Gramedia Pustaka Utama.
- [11] Widianingsih, M. (2017). Aktivitas antioksidan ekstrak metanol buah naga merah (Hylocereus polyrhizus (FAC Weber) Britton & Rose) hasil maserasi dan dipekarkan dengan kering angin. Jurnal Wiyata: Penelitian Sains dan Kesehatan, 3(2), 146-156.
- [12] Khoo, H. E., Azlan, A., Tang, S. T., & Lim, S. M. (2017). Anthocyanidins and anthocyanists colored pigments as food, pharmaceutical ingredients, and the potential health benefits. Food & Nutrition Research, 61(1), 1361779. doi:10.1080/16546628.2017.1361779.
- [13] Suyatma NE. (2009). Analisis Warna. Available online: http://slideplayer.info/slide/3239849/. (accessed on 14 April 2018).
- [14] Zahnfabrik (2017). Vita Easyshade. Available online: https://www.vita-zahnfabrik.com/cn/VITA-Easyshade-26934,27568.html (accessed on 17 April 2018).
- [15] Rianti, D., & Munadziroh, E. (2000). Perubahan Warna Resin Akrilik untuk Basis Gigi Tiruan dan Mahkota Jaket Akibat Jus Apel. Journal of Dentistry Indonesia, 7(3), 650-654.







DATA HASIL PENGAMATAN WARNA PLAT RESIN AKRILIK HEAT CURED

PRE-PENELITIAN CHROME

KONTROL	PERLAKUAN
-6.2	-7
-6.1	-5.4
-5.4	-5.4
-6.2	-6.2
-5.4	-6.6
-6.2	-6.5
-5.6	-5.3
-6.4	-7.7
-6.3	-6.1
-6.6	-5.5
-6.7	-5.7
-6	-5.6
-6.8	-6.3
-5.2	-6.6
-6.8	-6.2
-7.2	-6.8

POST PENELITIAN 7 HARI CHROME

KONTROL	PERLAKUAN
2.8	2.3
1.3	2.2
1.4	2.1
2	2.1
1.8	2.2
2.9	2.3
2.3	1.8
2	2.6
2	2.8
2.2	2.4
2.3	2.2
2.8	2.1
2	2.8
1.8	2.7
1.4	2.2
2.9	2.1

POST PENELITIAN 14 HARICHROME

KONTROL	PERLAKUAN
2.8	2.9
2.2	2.9
1.7	2.6
2.2	2.5
2	2.8
2.1	2.6
2.3	2.4
2.4	3.4
2.3	3.3
2.2	2.8
2.3	2.8
2.1	2.9
2.8	3.4
1.8	3.3
2.2	2.8
2.6	2.9

PRE-PENELITIAN VALUE

KONTROL	PERLAKUAN
-12.8	-13.6
-13.7	-15.1
-15.3	-13.1
-13.9	-15.6
-15.6	-13.3
-14.1	-15.5
-15.6	-15.9
-14.2	-13.2
-13.7	-14.5
-14.6	-13.3
-13.7	-15.3
-13.3	-15.7
-13.1	-14.1
-15.2	-13.3
-12.2	-13.4
-13.2	-15.6

POST PENELITIAN 14 HARIVALUE

KONTROL	PERLAKUAN
-25.1	-28.8
-27.1	-29.9
-25	-30.8
-27.7	-29.7
-24.7	-28.3
-25.6	-29.7
-24.7	-28.7
-25	-29.2
-26.1	-28.2
-26.5	-29.4
-27.3	-28.8
-25.2	-30.2
-27.4	-29.6
-24.6	-28.4
-25.7	-28.8
-24.8	-29.6

POST PENELITIAN 7 HARIVALUE

KONTROL	PERLAKUAN
-23.5	-22.4
-23	-23.1
-21.8	-23.3
-23.5	-22.3
-23.3	-21.1
-23.7	-22.2
-22.9	-22.3
-22.4	-23.1
-23.3	-22.5
-25.4	-23
-23.6	-23.1
-22.8	-23.2
-23.9	-22.4
-23.4	-21.3
-22.8	-22.4
-25.3	-22.2

LAMPIRAN 3

UJI STATISTIK *FRIEDMAN*

Warna Chrome

Ranks

	Mean Rank
Aquades sebelum perendaman	1.00
Aquades perendaman 7 hari	2.34
Aquades perendaman 14 hari	2.66

Test Statistics^a

16
6.881
2
.000

Interpretasi:

1. Didapatkan p = 0,000 (p <0,050) artinya terdapat perbedaan yang signifikan pada warna chrome kelompok control

Warna Chrome

Ranks

	Mean Rank
Larutan jus buah naga merah sebelum perendaman	1.00
Larutan jus buah naga merah perendaman 7 hari	2.00
Larutan jus buah naga merah perendaman 14 hari	3.00

Test Statistics^a

Ν	16
Chi-Square	32.000
Df	2
Asymp. Sig.	.000

Interpretasi:

1. Didapatkan p = 0,000 (p <0,050) artinya terdapat perbedaan yang signifikan pada warna chrome kelompok perlakuan

Warna Value

Ranks

	Mean Rank
Aquades sebelum perendaman	3.00
Aquades perendaman 7 hari	1.94
Aquades perendaman 14 hari	1.06

Test Statistics^a

N	16
Chi-Square	30.125
Df	2
Asymp. Sig.	.000

Interpretasi:

1. Didapatkan p = 0,000 (p < 0,050) artinya terdapat perbedaan yang signifikan pada warna value kelompok kontrol

Warna Value

Warna Value

Ranks

	Mean Rank
Larutan jus buah naga merah sebelum perendaman	3.00
Larutan jus buah naga merah perendaman 7 hari	2.00
Larutan jus buah naga merah perendaman 14 hari	1.00

Test Statistics^a

N	16
Chi-Square	32.000
Df	2
Asymp. Sig.	.000

Interpretasi:

1. Didapatkan p = 0,000 (p < 0,050) artinya terdapat perbedaan yang signifikan pada warna value kelompok kontrol

LAMPIRAN 4

UJI STATISTIK MANN WHITNEY

Warna Chrome

Sebelum Perendaman

Test Statistics^a

	Aquades dan Larutan jus buah naga merah
Mann-Whitney U	124.500
Wilcoxon W	260.500
z	132
Asymp. Sig. (2-tailed)	.895
Exact Sig. [2*(1-tailed Sig.)]	.897 ^b

Perendaman 7 hari

Test Statistics^a

	Aquades dan Larutan jus buah naga merah
Mann-Whitney U	94.000
Wilcoxon W	230.000
Z	-1.289
Asymp. Sig. (2-tailed)	.197
Exact Sig. [2*(1-tailed Sig.)]	.210 ^b

Perendaman 14 hari

Test Statistics^a

	Aquades dan Larutan jus buah naga merah
Mann-Whitney U	15.500
Wilcoxon W	151.500
Z	-4.266
Asymp. Sig. (2-tailed)	.000
Exact Sig. [2*(1-tailed Sig.)]	.000 ^b

Interpretasi:

- 1. Didapatkan p = 0,895 (p >0,050) artinya **tidak terdapat** perbedaan yang signifikan pada warna chrome kelompok kontrol dan larutan jus buah naga merah
- 2. Didapatkan p = 0,197(p >0,050) artinya **tidak terdapat** perbedaan yang signifikan pada warna chrome kelompok kontrol dan larutan jus buah naga merah
- 3. Didapatkan p = 0,000 (p >0,050) artinya **terdapat** perbedaan yang signifikan pada warna chrome kelompok control dan larutan jus buah naga merah

Warna Value

Sebelum Perendaman

Test Statistics^a

	Aquades dan Larutan jus buah naga merah sebelum perendaman
Mann-Whitney U	104.500
Wilcoxon W	240.500
Z	888
Asymp. Sig. (2-tailed)	.375
Exact Sig. [2*(1-tailed Sig.)]	.381 ^b

Perendaman 7 hari

Test Statistics^a

	Aquades dan Larutan jus buah naga merah
Mann-Whitney U	47.000
Wilcoxon W	183.000
z	-3.059
Asymp. Sig. (2-tailed)	.002
Exact Sig. [2*(1-tailed Sig.)]	.002b

Perendaman 14 hari

Test Statistics^a

	Aquades dan Larutan jus buah naga merah
Mann-Whitney U	.000
Wilcoxon W	136.000
Z	-4.828
Asymp. Sig. (2-tailed)	.000
Exact Sig. [2*(1-tailed Sig.)]	.000 ^b

Interpretasi:

- 1. Didapatkan p = 0,375 (p >0,050) artinya **tidak terdapat** perbedaan yang signifikan pada warna value kelompok kontrol dan larutan jus buah naga merah
- 2. Didapatkan p = 0,002 (p <0,050) artinya **terdapat** perbedaan yang signifikan pada warna value kelompok control dan larutan jus buah naga merah
- 3. Didapatkan p = 0,000 (p > 0,050) artinya **terdapat** perbedaan yang signifikan pada warna value kelompok kontrol dan larutan jus buah naga merah

Fwd: Accepted for publication

Kotak Masuk



Fransiska Nuning Kusmawati

15.43 (8 menit yang lalu)

kepada saya

Inggris Indonesia

Terjemahkan pesan

Nonaktifkan untuk: Inggris

----- Forwarded message ------

Dari: Fransiska Nuning Kusmawati < nuningphynx@gmail.com >

Date: Jum, 20 Sep 2019 pukul 13.09 Subject: Re: Accepted for publication

To: Andreas Demopoulos < demopoulos@scimedjournal.org>

Thanks, I have received it.

Pada tanggal Jum, 20 Sep 2019 12.37, Andreas Demopoulos demopoulos@scimedjournal.org menulis:

Dear Dr. Fransiska Nuning Kusmawati,

Greetings!

We are glad to inform you that your manuscript entitled "Effect of Red Dragon Fruit Juice on Acrylic Resin Color" has been accepted in the Scimed Journal Vol 1, No 3 (2019) by our reviewers and editors for publication without any further revisions.

We'll soon send you the final pdf version of your manuscript for your review and corrections before publication.

Please acknowledge the receipt of this email.

Thanks & Regards

Dr. Andreas Demopoulos Technical Editor