

AMJ

by Sinta Amj

Submission date: 13-Apr-2023 09:26AM (UTC+0700)

Submission ID: 2063074563

File name: AMJ_Dec_2022.pdf (253.6K)

Word count: 4350

Character count: 23578

Antibacterial Potential of *Stevia rebaudiana* Bertoni Leaf Extract on The Growth of *Enterococcus faecalis* Bacteria

Sinta Deviyanti¹, Silva Abraham², Burhanuddin Daeng Pasiga³

Oral Biology Department, Faculty of Dentistry, Prof.Dr.Moestopo University (B), Indonesia^{1,2}
Dental Public Health Department, Faculty of Dentistry, Prof.Dr.Moestopo University (B), Indonesia³



Keywords:

Stevia rebaudiana Bertoni,
Enterococcus faecalis, antibacterial,
irrigation solution, dental root canal

ABSTRACT

Enterococcus faecalis bacteria often found in cases of root canal treatment failure. Information on the antibacterial potential of *Stevia rebaudiana* Bertoni herbs against *E.faecalis*, is still very limited. To determine the antibacterial potential of *Stevia rebaudiana* Bertoni leaf extract against *Enterococcus faecalis*. In vitro laboratory experimental research with post test only design with control group. *Stevia* leaf extraction using maceration method. The antibacterial activity of *Stevia* leaf extract against *Enterococcus faecalis* was tested by agar well diffusion method. A total of 50 µl of *Stevia* leaf extracts in different concentration with positive control NaOCL 2.5% and negative control distilled water, were put into the wells on Nutrient agar plates that have been inoculated with *E.faecalis*. Incubate at 37°C for 24 hours. Zone of inhibition (ZOI) was measured using a caliper. The ZOI of all test concentrations of *Stevia* leaf extract were varied, with maximum diameter of 23.07 mm and a minimum of 10.34 mm, significantly different ($P<0.05$) from the negative control. The growth inhibition of *faecalis* bacteria by 100% and 50% *Stevia* leaf extracts was not significantly different ($P>0.05$) compared to the positive control while 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia* leaf extracts were significantly different ($P<0.05$) compared to the positive control. All test concentrations of *Stevia* leaf extract have the potential to inhibit the growth of *Enterococcus faecalis* bacteria. *Stevia* leaf extracts of 100% and 50% have the ability to inhibit *E.faecalis* bacteria which is equivalent to NaOCl 2.5%.



28

This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License.

1. Introduction

Enterococcus faecalis is a facultative anaerobic gram-positive bacterium that is often found in dental pulp infections and 80-90% are found in cases of infection due to failure of root canal treatment (secondary endodontic infection) [1], [2]. *Enterococcus faecalis* is also known to be the most virulent bacteria and resistant to antibiotics (such as penicillin G, vancomycin) and antiseptics. This bacterium is difficult to

eradicate because it is able to survive (persistence) in dentinal tubules in the root canal as an area with limited access to antibiotics and antiseptics.¹ These bacteria can also survive in the root canals of obturated teeth for at least 6 months even without the availability of nutrients from outside [3]. The ability of these bacteria to survive in the root canal system and cause secondary endodontic infections because they have the ability to adhere to dentin and invade dentinal tubules, are able to live in high pH and low pH environments, and are able to form biofilms, which contribute to resistance and persistence after antimicrobial procedures in the root canal [1], [4]. The pathogenicity of these bacteria ranges from causing life-threatening disease in compromised individuals to less severe conditions such as infection in the root canals of obturated teeth with chronic apical periodontitis.⁴ The virulence factors of this bacterium most associated with endodontic infection and peri-radicular inflammatory response are aggregation substance, surface adhesion, sex pheromones, lipoteichoic acid, extracellular superoxide production, the lytic enzymes gelatinase and hyaluronidase and the toxin cytolysin [4].

Medicinal plants are becoming very popular for the treatment of different diseases all over the world. *Stevia rebaudiana* Bertoni is one of the herbs known to have more than 100 types of phytochemical content, has been used in ancient medical systems for antimicrobial, antifungal, anti-oxidative, antitumor, anti-hypertension, hepatoprotective, hypoglycemic, and antiviral activity. *Stevia rebaudiana* Bertoni is a species of plant from the Asteraceae family.

This plant is known as honey leaf or sweet leaf because it contains Steviol glycoside component as a natural non-caloric sweetener that causes a sweet taste in its leaves. This plant originates from the highland areas of northeastern Paraguay and currently the *Stevia* plant has been recognized in various countries including Indonesia [5].

Various studies have proven that *Stevia rebaudiana* Bertoni leaf extract has antibacterial activity against various types of bacteria and fungi, such as *Streptococcus mutans*, *Streptococcus sobrinus*, *Streptococcus salivarius*, *Streptococcus mitis*, *Lactobacillus acidophilus*, *Lactobacillus casei*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Pseudomonas aeruginosa* and *Candida albicans* [6- 8]. However, information on research results related to the antibacterial activity of *Stevia rebaudiana* Bertoni leaf extract against the growth of *Enterococcus faecalis* bacteria, is still very limited until now. The purpose of this study was to determine the antibacterial potential of *Stevia rebaudiana* Bertoni leaf extract against the growth of *Enterococcus faecalis* bacteria.

2. Materials and Methods

2.1 Determination assay of *Stevia rebaudiana* Bertoni plant

Determination tests of *Stevia rebaudiana* Bertoni plants purchased from plantations in Ciputri village, Pacet sub-district, Cianjur district in West Java, were carried out at Herbarium Bogoriense, Botany Division-BRIN Biological Research Center in Cibinong, Bogor.

2.2 Phytochemical screening of *Stevia rebaudiana* Bertoni leaf extract

Detection of terpenoids (*Salkowski's test*) is done by adding 2 ml of chloroform (CHCl₃) and 3ml of concentrated sulfuric acid (H₂SO₄) to 1ml of *Stevia* leaf extract solution. If a reddish brown color is formed, it indicates the content of terpenoid phytochemicals [9], [10]. Flavonoid detection is done by adding 5 ml of liquid ammonia (NH₃) and 1 ml of concentrated sulfuric acid (H₂SO₄) into 1 ml of *Stevia* leaf extract solution. If a yellow color is formed, it indicates the content of flavonoid phytochemicals.

¹⁰Steroid detection is done by adding 1 ml of anhydrous acetic acid (CH₃ COOH) and 1 ml of concentrated

sulfuric acid (H₂SO₄) into 1 ml of *Stevia* leaf extract solution. If a green color is formed, it indicates the content of steroid phytochemicals [10]. Detection of tannin is done by adding 1 ml of hydrogen chloride (HCL) and 1ml of Dragendorff's reagent (potassium bismuth solution) into 1 ml of *Stevia* leaf extract solution. If a reddish brown or reddish orange color is formed, it indicates the content of alkaloid phytochemicals [10], [11]. Alkaloid detection is done by adding 1 ml of ferric chloride (FeCL₃) into 1 ml of *Stevia* leaf extract solution. When formed green or purplish green color, indicating the content of alkaloid phytochemicals [9], [10]. Detection of saponins (foam test) is done by putting 0.5 g of *Stevia* leaf extract into 2 ml of water and then shaken. If foam is formed that lasts for 10 minutes, it indicates the content of saponin phytochemicals [12].

2.3 Preparation of extraction of *Stevia rebaudiana* Bertoni leaf with maseration methods

Dry leaf samples of *Stevia rebaudiana* Bertoni were pulverized with a blender and filtered using a mesh (100 mesh) to obtain 150 gr of simplisia which was poured into 3 Duran bottles so that each bottle contained 50 gr of *Stevia* leaf simplisia. The preparation of *Stevia* leaf extract was carried out through the maceration method by means of each Duran bottle then filled with 96% ethanol solvent as much as 400 ml and soaking for 3x24 hours at room temperature (25⁰ C). The maceration solution was shaken periodically every 8 hours for 15 minutes each day. Furthermore, filtering was carried out with a glass funnel lined with Whatman paper and evaporation of the macerated solution using a rotary evaporator at a temperature of 72⁰ C with a rotation speed of 60 rpm and a vacuum pump pressure of 900 mBar until a thick extract of *Stevia* leaves was obtained [12- 14]. *Stevia* leaf extract was stored in a -20⁰ C freezer before being used for testing.

2.4 Preparation of *E.faecalis* culture bacteria

E. faecalis strain ATCC 29212 was cultured by growing 20 µl of bacteria on 5 ml of Nutrient broth medium and incubated for 24 hours at 37°C. Optical Density (OD) measurement of the bacterial suspension was carried out using a microplate reader at a wavelength of 600 nm and equalized the amount of bacterial density according to the Mc Farland standard of 1.5x10⁸ CFU/ml (Mc Farland 0.5).

2.5 Preparation of *Stevia rebaudiana* Bertoni leaf extract

The preparation of 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia* leaf extract solution was done through serial dilution method in eppendorf tubes. A total of 1 g *Stevia* leaf extract was dissolved in 1 ml sterile distilled water to obtain 100% *Stevia* leaf extract solution in tube 1. The percentage of 50% *Stevia* leaf extract solution was made by transferring 500 µl of extract in tube 1 to tube 2 containing 500 µl sterile distilled water. The solution of 25% *Stevia* leaf extract was made by transferring 500 µl of extract in tube 2 to tube 3 containing 500 µl of sterile distilled water. The serial dilution procedure was continued until a solution of 1.625% *Stevia* leaf extract was obtained.

2.6 Detection of antibacterial activity of *Stevia rebaudiana* Bertoni leaf extract against *E.faecalis*

The *E. faecalis* antibacterial activity of *Stevia* leaf extract was tested using agar well diffusion method [15]. Sterile Nutrient agar media on 7 Petri dishes were prepared and incubated for 24 hours at 37⁰ C. A total of 20 µl of fresh suspension of *E. faecalis* bacteria that has been equalized with Mc Farland 0.5 standard was streaked onto the surface of solid Nutrient Agar media in Petri dishes. Five wells in each Petri dish were made using a 6 mm diameter cork borer. A total of 3 wells in each petri dish (triplo), each filled with 1 type of specific concentration of *Stevia* leaf extract solution as much as 50 µl. The other two wells were each filled with positive control (NaOCL 2.5%) and negative control (sterile distilled water) by 50 µl in each petri dish. A total of 7 Petri dishes were then incubated under anaerobic conditions at 37⁰ C for 24 hour. The mean ZOI formed was measured using a caliper in 3 directions (vertical, horizontal, diagonal) for each sample [16].

2.7 Statistical analysis

Statistical analysis was performed by Kruskal Wallis and post hoc Mann-Whitney test using the SPSS software version 26.

3. Results

Stevia rebaudiana Bertoni Plant Determination Test Results

The results of the determination test stated that the tested plant is the species *Stevia rebaudiana* Bertoni from the Asteraceae family.

3.1 Results of Maceration Extraction of *Stevia rebaudiana* Bertoni Leaves

The results of the extraction of 150 grams of *Stevia rebaudiana* Bertoni leaf simplisia by maceration method for 3x24 hours using 1200 ml of 96% ethanol solvent, followed by filtration then the filtrate is evaporated with a rotary vacuum evaporator for 1.5 hours at a temperature of 72⁰ C can produce a thick extract weighing 22.47 grams.

3.2 Phytochemical Screening Result of *Stevia rebaudiana* Bertoni Leaf Extract

Phytochemical screening results of *Stevia rebaudiana* Bertoni leaf extract in this study showed the presence of bioactive compounds terpenoids, flavonoids, steroids, tannins, alkaloids and saponins (Table 1).

Table 1. Phytochemical screening results of ethanol extract of *Stevia rebaudiana* Bertoni leaves

PHYTOCHEMICAL TEST	ETHANOLIC EXTRACT OF <i>Stevia rebaudiana</i> Bertoni LEAVES	INDICATOR
Terpenoids	+	A reddish-brown color is formed,
Flavonoids	+	Yellow color formed
Steroids	+	Green color formed
Tannins	+	Forms a green or purplish green color
Alkaloids	+	Reddish brown or reddishorange color is formed
Saponins	+	Forms a foam that lasts for 10 minutes

3.3 Antibacterial activity of *Stevia* leaf extract irrigation solution against *E. faecalis*:

The antibacterial activity test results of *Stevia rebaudiana* Bertoni leaf extract concentrations of 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and 1.56% and the control group against *E. faecalis* bacteria by agar well diffusion method (Figure 1a, 1b, 1c, 1d, 1e, 1f, 1g, 1h).




Figure 1 Zone of inhibition (ZOI) of *E. faecalis* bacteria by *Stevia* leaf extract
(A) 100%, K+ and K-, (B) 50%, K+ and K-, (C) 25%, K+ and K-, (D) 12.5%, K+ and K-,
(E) 6.25%, K+ and K-, (F) 3.125%, K+ and K-, (G) 1.56%, K+ and K-

The mean ZOI of *E. faecalis* bacteria (mm) in the *Stevia rebaudiana* Bertoni leaf extract solution group in various test concentrations as well as the control group can be seen in table 2.

Table 2. The mean ZOI of *E. faecalis* bacteria (mm) in the *Stevia* leaf extract groups in various test concentrations as well as the control group

		Stevia 100%	Stevia 50%	Stevia 25%	Stevia 12.5%	Stevia 6.25%	Stevia 3.125%	Stevia 1.56%	Control +	Control -
N	Valid	3	3	3	3	3	3	3	7	7
	Missing	0	0	0	0	0	0	0	0	0
Mean		23.0667	22.8433	21.0667	17.7900	16.4433	13.5567	10.3433	25.8714	.0000
Std. Deviation		1.51767	.80600	2.72015	.91995	1.83538	.87552	1.02788	2.04834	.00000
Minimum		21.70	22.10	18.13	17.00	14.63	12.63	9.73	22.40	.00
Maximum		24.70	23.70	23.50	18.80	18.30	14.37	11.53	29.00	.00

The diagram of mean ZOI of *E. faecalis* bacteria (mm) formed in the treatment groups of *Stevia rebaudiana* Bertoni leaf extract in various concentrations tested as well as the positive control group (NaOCL 2.5%) and negative control (aquadest) can be seen in figure 2.




Figure 2. Diagram of mean ZOI of *Enterococcus faecalis* bacteria (CFU/ml) in 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia rebaudiana* Bertoni leaf extract groups and K+ (NaOCL 2.5%) and K- (aquadest) groups.

The results of the Shapiro-Wilk normality test ($n < 50$) for 100%, 50%, 25%, 12.5%, 6.25% and 3.125% *Stevia rebaudiana* Bertoni leaf extract showed that the data were normally distributed ($p > 0.05$), except for 1.56% *Stevia rebaudiana* Bertoni leaf extract, positive control and negative control ($p < 0.05$). The results of the Kruskal-Wallis test of various concentrations of *Stevia* leaf extract and the control group on the mean ZOI of *E. faecalis* bacteria, obtained a value of $p = 0.000$ ($p < 0.05$), indicating that there is a difference in the mean ZOI of *E. faecalis* bacteria at various concentrations of *Stevia* leaf extract and control group so that it is continued with the Mann-Whitney post hoc test at the 95% confidence level. The results of the Mann-Whitney post hoc test (table 3) showed that the 100% and 50% *Stevia* leaf extract solution groups did not show a significant difference in mean ZOI of *E. faecalis* bacteria with the positive control group ($p > 0.05$). The 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia* leaf extract solution groups showed significant differences in mean ZOI of *E. faecalis* bacteria with the positive control group ($p < 0.05$). The 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia* leaf extract solution groups showed significant mean differences in ZOI of *E. faecalis* bacteria with the negative control group ($p < 0.05$).

Table 3. Results of Post Hoc Mann-Whitney ZOI Test of *E. faecalis* Bacteria Growth

	Stevia 100%	Stevia 50%	Stevia 25%	Stevia 12.5%	Stevia 6.25%	Stevia 3.125%	Stevia 1.56%	Control +	Control -
Stevia 100%		.827	.275	.050	.050	.050	.050	.051	.003
Stevia 50%	.827		.275	.050	.050	.050	.050	.051	.003
Stevia 25%	.275	.275		.127	.127	.050	.050	.029	.003
Stevia 12.5%	.050	.050	.127		.275	.050	.050	.016	.003
Stevia 6.25%	.050	.050	.127	.275		.050	.050	.016	.003
Stevia 3.125%	.050	.050	.050	.050	.050		.050	.016	.003
Stevia 1.56%	.050	.050	.050	.050	.050	.050		.016	.003
Control +	.051	.051	.029	.016	.016	.016	.016		.001
Control -	.003	.003	.003	.003	.003	.003	.003	.001	

Notes: Significant at $p < 0.05$

4. Discussion

Determination of *Stevia rebaudiana* Bertoni plants was carried out to ensure the correct identity of plant samples to be used in this study [17]. The results of the *Stevia* plant sample determination test used in this study, have shown the suitability of the taxonomic hierarchy of the *Stevia rebaudiana* Bertoni plant which states that the plant tested is the species *Stevia rebaudiana* Bertoni from the Asteraceae family [5].

The antibacterial activity test of *Stevia rebaudiana* Bertoni leaf extract against *E. faecalis* ATCC 29212 bacteria in this study was carried out using agar well diffusion method because it has the advantage of being able to accommodate more antibacterial agents tested compared to disc diffusion method so that it is expected that the inhibition zone formed is larger [18]. All variations of concentration of *Stevia rebaudiana* Bertoni leaf ethanol extract tested in this study, proved to be able to inhibit the growth of *E. faecalis* bacteria which was significantly different when compared to the negative control group ($p < 0.05$). Mean ZOI appeared to be greater in line with the increase in concentration with the largest and smallest mean ZOI values of *Stevia rebaudiana* Bertoni leaf extract solutions of 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and 1.56% were 23.07 mm, 22.84 mm, 21.07 mm, 17.79 mm, 16.44 mm, 13.56 mm and 10.34 mm respectively. The mean ZOI of the positive control (NaOCL 2.5%) was 25.87 mm while the negative control (aquadest) showed ZOI. The ability of all concentration variations of *Stevia rebaudiana* Bertoni leaf extract solution tested in this study to inhibit the growth of *E. faecalis* bacteria is possible because it contains various bioactive compounds with antibacterial activity. Phytochemical testing in this study proved that *Stevia rebaudiana* Bertoni leaf extract contains bioactive compounds (phytochemicals) including terpenoids, flavonoids, steroids, tannins, alkaloids and saponins. The content of phytochemical compounds detected in *Stevia rebaudiana* Bertoni leaf extract in this study seems to be in line with the results of several other researchers. Research by [19] has reported that the phytochemical content in ethanol and methanol extracts of *Stevia rebaudiana* Bertoni leaves are phenols, tannins, saponins, glycosides and flavonoids. Research by [8] has also proven the phytochemical content of alkaloids, steroids, tannins, saponins and flavonoids in *Stevia rebaudiana* Bertoni leaves. The antibacterial activity of tannin phytochemicals against *E. faecalis* occurs through the mechanism of inhibiting bacterial cell wall synthesis by forming irreversible complexes with proline-rich proteins. The antibacterial activity of saponin phytochemicals occurs through its ability to break down proteins and enzymes from bacterial cells. The antibacterial activity of terpenoid phytochemicals, occurs through the dissolution of bacterial cell walls. Furthermore, flavonoid phytochemicals play an effective role in forming complexes with soluble proteins and bacterial cell walls. Steroidal phytochemicals have the ability of bacterial activity through the mechanism of liposome breakdown [20]. The antibacterial activity of alkaloid phytochemicals, occurs through inhibition of bacterial cell wall synthesis and disruption of the formation of peptidoglycan components in bacterial cells resulting in failure of bacterial cell wall layer formation [21]. Mean ZOI of *E. faecalis* bacteria growth of *E. faecalis* bacteria from 100% and 50% *Stevia rebaudiana* Bertoni leaf extract solution which is not significantly different ($p > 0.05$) from the positive control group from the results of this study, indicating that both concentrations of *Stevia* leaf extract have antibacterial activity that is as good as the positive control (NaOCL 2.5%). The growth inhibition response category of *E. faecalis* bacteria from 100% and 50% *Stevia rebaudiana* Bertoni leaf extract is classified as very strong according to Davis W and Stout TR (1971) cited by [22]. The group of *Stevia rebaudiana* Bertoni leaf extract with concentration of 25%, 12.5%, 6.25%, 3.125% and 1.56% showed a difference in the mean ZOI of *E. faecalis* bacteria when compared to the positive control group ($p < 0.05$) with the response category of *E. faecalis* bacteria growth inhibition classified as strong according to Davis W and Stout TR (1971) cited by [22].

The ability of the entire *Stevia rebaudiana* Bertoni leaf extract in ethanol solvent to inhibit the growth of *E. faecalis* bacteria from the results of this study, seems to be in line with the research of Ghosh S *et al* cited by [6] states that the concentration of *Stevia rebaudiana* Bertoni leaf extract in ethanol solvent

concentration of 250 µg/ml and various other types of solvents, able to inhibit the growth of various bacteria and fungi including *E. faecalis* bacteria. The diameter of the inhibition zone of ethanol extract of *Stevia rebaudiana Bertoni* leaves against *E. faecalis* MTCC 2729 bacteria by disc diffusion method in ethanol, water, chloroform, cyclohexane, acetone and petroleum ether solvents from the results of the study in order are 5 mm, 5 mm, 6mm, 8.3 mm and 13 mm. The positive control (streptomycin) in the study produced an inhibition zone diameter of 18 mm.

The use of sodium hypochlorite (NaOCL) 2.5% as a positive control in this study because it has a very strong antimicrobial capacity [3], [23]. This is evidenced by the mean ZOI NaOCL 2.5% which is the greatest against the growth of *E. faecalis* bacteria compared to all groups of *Stevia rebaudiana Bertoni* leaf extract solutions and negative controls. The antimicrobial activity of NaOCL is based on high pH conditions as a strong base (pH> 11) which can disrupt the integrity of the cytoplasmic membrane of bacterial cells due to irreversible enzyme inhibition, cause biosynthetic changes in cellular metabolism and cause phospholipid degradation [23].

5. Conclusion

All *Stevia rebaudiana* leaf extract tested in this study were proven to have antibacterial potential of *Enterococcus faecalis* ATCC 29212 because they were able to form ZOI of bacterial growth.

Stevia rebaudiana Bertoni leaf extract 100% and 50% have a very strong inhibition response category with antibacterial potential to inhibit the growth of *Enterococcus faecalis* ATCC 29212 equivalent to NaOCL 2.5%.

Stevia rebaudiana Bertoni leaf extract 25%, 12.5%, 6.25%, 3.125% and 1.56% have a strong inhibitory response category and have antibacterial potential to inhibit the growth of *Enterococcus faecalis* ATCC 29212 which is significantly different from the positive control group.

The antibacterial ability of 25%, 12.5%, 6.25%, 3.125% and 1.56% *Stevia rebaudiana Bertoni* leaf extracts because they are proven to contain phytochemicals including terpenoids, flavonoids, steroids, tannins, alkaloids and saponins.

5.1 Advice

Further research is needed to test the effectiveness and toxicity of *Stevia rebaudiana Bertoni* leaf extract against *E. faecalis* bacteria so that its clinical utilization can be more optimal, among others, as an alternative solution for dental root canal irrigation.

6. References

- [1] Samaranayake L. Essential microbiology for dentistry 5th ed. Edinburgh: Elsevier. 2018:129,310-311.
- [2] Fouad AF. Endodontic microbiology. 2nd ed. Singapore: John Wiley & Sons. 2017:129-170.
- [3] Bjorndal L, Kirkevang LL, Whitworth J. Textbook of endodontology. 3rd ed. New Delhi: John Wiley & Sons Ltd. 2018. Pg: 231-236.
- [4] Borzini L, Condo R, Dominici PD, Casaglia A, Cerroni L. Root canal irrigation: chemical and plant extracts against *Enterococcus faecalis*. The Open Dentistry Journal. 2016;10:692-703.

- [5] Gunasena MDKM, Senarath RMUS, Senarath WTPSK. ¹³ A Review on Chemical Composition, Biosynthesis of Steviol Glycosides, Application, Cultivation, and Phytochemical Screening of *Stevia rebaudiana* (Bert.) Bertoni. *Journal of Pharmaceutical Research International*.2021;33(29B):85-104.
- [6] ¹⁷ Gamboa F, Chaves M. Antimicrobial Potential of Extract from *Stevia rebaudiana* Leaves Against Bacteria of Importance in Dental Caries. *Acta Odontologica Latinoamericana*. 2012;2(25):171-175.
- [7] Arambula MM, Alvarado MO, Sanchez ML. ¹ Antibacterial Activity of Extract of *Stevia rebaudiana* Bertoni Against *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Pseudomonas aeruginosa*. *Journal of Medicinal Plant Research*. 2017;11(25):414-418.
- [8] ¹² Siddique AB, Rahman SMM, Hossain MA, Rashid MA. Phytochemical Screening and Comparative antimicrobial potential of different extracts of *Stevia rebaudiana* Bertoni leaves. *Asian Pacific Journal of Tropical Disease*. 2014;4(4):275-280. [https://doi.org/10.1016/S2222-1808\(14\)60572-7](https://doi.org/10.1016/S2222-1808(14)60572-7)
- [9] ⁶ Pant DR, Pant ND, Saru DB, Yadav UN, Khanal DP. Phytochemical Screening and Study of Antioxidant, Antimicrobial, Antidiabetic, Anti-inflammatory and Analgesic Activities of Extracts From Stem Wood of *Pterocarpus marsupium* Roxburgh. *Journal of Intercultural Ethnopharmacology*.2017;
- [10] ¹⁰ Zivic N, Milosevic S, Dekic V, Dekic B, Ristic N, Ristic M, Sretic L. Phytochemical and Antioxidant Screening of Some Extracts of *Juniperus communis* L. *Czech Journal of Food Sciences*.2019; 37(5): 351-358.
- [11] ¹⁹ Hasan A, Akmal Z, Khan N. The phytochemical Screening and Antioxidants Potential of *Schoenoplectus triquetra* L.Palla. *Journal of Chemistry*.2020:1-8
- [12] ⁹ Pandey A, Tripathi S. Concept of Standardization, Extraction, and Phytochemical Screening Strategies for Herbal Drug. *Journal of Pharmacognosy and Phytochemistry*. 2014;2(5):115-119.
- [13] ¹⁶ Endarini LH. Pharmacognition and phytochemistry. Ministry of Health of the Republic of Indonesia-Health Human Resources Education Center.2016: 145.
- [14] ²⁶ Chandra Chandra A. Studi Awal Ekstraksi Batch Daun *Stevia rebaudiana* dengan variable Jenis Pelarut dan Temperatur Ekstraksi. *Pros Sem Nas Masy Biodiv Indon*.2015;1(1):114-115.
- [15] ¹⁴ Bolouri M, Sadiki M, Ibsouda SK. Methods for In Vitro Evaluating Antimicrobial Activity: A Review. *Journal of Pharmaceutical Analysis*. 2016;6:71-79.
- [16] ³ Mozartha M, Silvia P, Sujatmiko B. Comparison of Antibacterial activity of *Curcuma zedoaria* Extract and 2.5% Sodium Hypochlorite Irrigation Materials against *Enterococcus faecalis*. *Journal of Materials Dentistry*.2019;8(1):22-29.
- [17] ²⁵ Rumagit HM, Runtuwene MRJ, Sudewi S. Uji fitokimia dan Uji Aktivitas Antioksidan Dari Ekstrak Etanol Spons *Lamellodysidea herbacea*. *Pharmakon Jurnal Ilmiah Farmasi*.2015;4(3): 183-190.
- [18] ⁵ Khusuma A, Safitri Y, Yuniarni A, Rizki K. Uji Teknik Difusi Menggunakan Kertas Saring media Tampung Antibiotik Dengan *Escherichia coli* Sebagai Bakteri Uji. *Jurnal Kesehatan*

Prima.2019;13(2):151-155.

[19] Gupta E, Vajpayee G, Purwar S, Shakyawar S, Alok S, Shanthi A, Sundaram. ³⁴Phytochemical Screening and in-Vitro ³⁸Studies of Antioxidant and Antimicrobial Activity of Extracts of Dried Stevia rebaudiana Leaves. ⁴¹International Journal of Pharmaceutical Sciences and Research. 2017;8(8): 3354–3360. [https://doi.org/10.13040/IJPSR.0975-8232.8\(8\).3354-60](https://doi.org/10.13040/IJPSR.0975-8232.8(8).3354-60)

⁸[20] Mujeeb F, Bajpai P, Pathak N. Phytochemical evaluation, antimicrobial activity, and determination of bioactive components from leaves of aegle marmelos. BioMed Research International, 2014. <https://doi.org/10.1155/2014/497606>

¹[21] Hastuty A. Antibiofilm and antimicrobial activities of papaya (Carica papaya L.) and stevia (Stevia rebaudiana Bertoni) leaf extracts against three biofilm-forming bacteria. Journal of Microbial Systematics and Biotechnology, 2019;1(1): 19-29. <https://doi.org/10.37604/jmsb.v1i1.18>

²⁹[22] Widyani KAM, Wawengkang DS, Rumondor EM. Potensi Ekstrak dan Fraksi Spons Mycale vansoesti dari Perairan Manado Tua Terhadap Pertumbuhan Bakteri Staphylococcus aureus dan Escherichia coli. Pharmacon.2022;11(3): 1583-1590.

[23] Berman LH, Hargreaves KM, Rostein I. Cohen's Pathways of the Pulp. 12th ed. St. Louis: Elsevier 2021:272-281.

23%

SIMILARITY INDEX

%

INTERNET SOURCES

23%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

Aerma Hastuty. "Antibiofilm and antimicrobial activities of papaya (*Carica papaya* L.) and stevia (*Stevia rebaudiana* Bertoni) leaf extracts against three biofilm-forming bacteria", *Journal of Microbial Systematics and Biotechnology*, 2019

Publication

2%

2

Letizia Borzini, Roberta Condò, Paolo De Dominicis, Adriano Casaglia, Loredana Cerroni. "Root Canal Irrigation: Chemical Agents and Plant Extracts Against *Enterococcus faecalis*", *The Open Dentistry Journal*, 2016

Publication

2%

3

Tessya Nitulo Permatasari Lahagu, Huynh Diem An, Cindy Denhara Wijaya, Mellisa Sim. "Efficacy of *Cyombopogon Citratus* Extract Against *Enterococcus Faecalis*", *Biomedical Journal of Indonesia*, 2021

Publication

2%

4

Djéssica Tatiane Raspe, Camila da Silva, Silvio Cláudio da Costa. "Compounds from Stevia rebaudiana Bertoni leaves: An overview of non-conventional extraction methods and challenges", Food Bioscience, 2022

Publication

1 %

5

G N Mediarman, Sumardianto, P H Riyadi, L Rianingsih, L Purnamayati. "Potentials of CaO powder result of calcination from green shells (*Perna viridis*), scallops (*Placuna placenta*), and blood clams (*Anadara granosa*) as antibacterial agent", IOP Conference Series: Earth and Environmental Science, 2021

Publication

1 %

6

N Rashid, AA Paul, S Islam, SA Sajib, K Nasirujjaman, KMF Hoque, MA Reza. "Studies on antioxidant potential, phytochemical properties and toxicity of four popular medicinal plants of Bangladesh", Journal of Bio-Science, 2018

Publication

1 %

7

Aldelany Ramalho Freire, Deborah Ellen Wanderley Gomes Freire, Simone Alves de Sousa, Eliane Medeiros Serpa et al. "Antibacterial and Solubility Analysis of Experimental Phytotherapeutic Paste for Endodontic Treatment of Primary Teeth",

1 %

Pesquisa Brasileira em Odontopediatria e Clínica Integrada, 2021

Publication

8

S. Abirami, J. Jabesta, R. Emilin Renitta, D. Alex Anand, Antony V. Samrot. "ANTIMICROBIAL ACTIVITY OF FLOWER EXTRACTS AGAINST WOUND PATHOGENS", Current Research in Green and Sustainable Chemistry, 2021

Publication

1 %

9

W. F. Briggs, H. O. Stanley, G. C. Okpokwasili, O. M. Immanuel, C. J. Ugboma. "Biocorrosion Inhibitory Potential of Aqueous Extract of Phyllanthus amarus against Acid Producing Bacteria", Journal of Advances in Biology & Biotechnology, 2019

Publication

1 %

10

Bachiri Fatiha, Benouaklil Fatouma, Hamaidi-Chergui Fella, Saidi Fairouz. " Antibacterial Activity of Essential Oil from Ripe Berries of L. Against Multi-drug Resistant Pathogens Responsible for Skin Infections in Burn Patients ", Journal of Essential Oil Bearing Plants, 2022

Publication

1 %

11

IFMBE Proceedings, 2016.

Publication

1 %

12

P. A. S. Theophilus, M. J. Victoria, K. M. Socarras, K. R. Filush, K. Gupta, D. F. Luecke,

1 %

E. Sapi. " Effectiveness of whole leaf extract against the various morphological forms of ", European Journal of Microbiology and Immunology, 2015

Publication

13

Yang Wang, Xiang Luo, Li Chen, Abdullateef Taiye Mustapha, Xiaojie Yu, Cunshan Zhou, Clinton Emeka Okonkwo. "Natural and low - caloric rebaudioside A as a substitute for dietary sugars: A comprehensive review", Comprehensive Reviews in Food Science and Food Safety, 2022

Publication

1 %

14

"Synthesis and physicochemical studies of fluconazole ionic liquids", Journal of Applied Pharmaceutical Science, 2017

Publication

<1 %

15

Asmaliyah, E E W Hadi, E Novriyanti. "Potential of medang reso (Cinnamomum parthenoxylon) as raw material source for antidiabetic drugs", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

16

Burhanuddin Daeng Pasiga, Arifurrahman Burhanuddin. "Emergency Intervention Need for Community Dental Services (Pathfinder Survey in North Mamuju Regency, West

<1 %

17

Pamela Nair Silva-Holguín, Simón Yobanny
Reyes-López. "Alumina-Hydroxyapatite-Silver
Spheres With Antibacterial Activity", Dose-
Response, 2021

Publication

<1 %

18

"Natural Oral Care in Dental Therapy", Wiley,
2020

Publication

<1 %

19

Priyanka Danai, Varsha Pandey, Teena
Agrawal. "In vitro antioxidant potential and
antimicrobial activity of leaves and stem
extracts of Anogeissus pendula Edgew.", Plant
Science Today, 2021

Publication

<1 %

20

Farina Mujeeb, Preeti Bajpai, Neelam Pathak.
" Phytochemical Evaluation, Antimicrobial
Activity, and Determination of Bioactive
Components from Leaves of ", BioMed
Research International, 2014

Publication

<1 %

21

Zahrasadat Madani, Majid Sales, Ali Akbar
Moghadamnia, Sohrab Kazemi, Fariba
Asgharpour. "Propolis nanoparticle enhances
antimicrobial efficacy against Enterococcus

<1 %

22

"MOLECULAR AND MICROBIOLOGICAL ASSESSMENT OF STEVIA (*Stevia rebaudiana* BERTONI) LEAF EXTRACTS FOR ANTI-BACTERIAL AND ANTI-MUTAGENIC ACTIVITIES", Zagazig Journal of Agricultural Research, 2017

Publication

23

N Kamalakkannan, P Stanely Mainzen Prince. "Antihyperlipidaemic effect of *Aegle marmelos* fruit extract in streptozotocin-induced diabetes in rats", Journal of the Science of Food and Agriculture, 2005

Publication

24

Balakrishnan Arumugam, Arunambiga Subramaniam, Praveena Alagaraj. "Stevia as a Natural Sweetener: A Review", Cardiovascular & Hematological Agents in Medicinal Chemistry, 2020

Publication

25

Eka Cahya Nugraha, Tri Mulyowati, Rinda Binugraheni. "Uji Aktivitas Larvasida Ekstrak Etanolik Daun Serai Wangi (*Cymbopogon nardus* L.) terhadap Larva *Culex* sp. Instar III", Biomedika, 2019

Publication

<1 %

<1 %

<1 %

<1 %

26

Silfiana Nisa Permatasari, Umarudin ..
"Determinasi dan Analisa Proksimat Daun
Benalu pada Pohon Mangga Arum Manis di
Ketintang Madya Surabaya", Journal of
Pharmacy and Science, 2019

Publication

<1 %

27

Roberto Lemus-Mondaca, Antonio Vega-
Gálvez, Liliana Zura-Bravo, Kong Ah-Hen.
"Stevia rebaudiana Bertoni, source of a high-
potency natural sweetener: A comprehensive
review on the biochemical, nutritional and
functional aspects", Food Chemistry, 2012

Publication

<1 %

28

Shiva Khatibi, Zahra Farzaneh Taban, Amaneh
Mohammadi Roushandeh. "In Vitro Evaluation
of Cytotoxic and Antiproliferative Effects of
Portulaca oleracea Ethanolic Extracton on
HeLa Cell Line", Gene, Cell and Tissue, 2016

Publication

<1 %

29

Brigita Michelle Luntungan, Defny S.
Wewengkang, Erladys Rumondor. "UJI
AKTIVITAS ANTIBAKTERI EKSTRAK DAN FRAKSI
SPONS Mycale Vansoesti DARI PERAIRAN
PULAU MANTEHAGE MINAHASA UTARA
TERHADAP PERTUMBUHAN BAKTERI
Escherichia coli DAN Staphylococcus aureus",
PHARMACON, 2021

Publication

<1 %

30

Wendkouni Leila Marie Esther Belem-Kabré, Vincent Ouédraogo, Bagora Bayala, Alimata Bancé et al. "Anti-Biofilm, Anti-Quorum Sensing, and Anti-Proliferative Activities of Methanolic and Aqueous Roots Extracts of *Carica papaya* L. and *Cocos nucifera* L.", *Advances in Microbiology*, 2023

Publication

<1 %

31

Andreas Romulo, Ervizal A. M. Zuhud, Johana Rondevaldova, Ladislav Kokoska. " Screening of antimicrobial activity of plants used in traditional Indonesian medicine ", *Pharmaceutical Biology*, 2018

Publication

<1 %

32

Ferrazzano, Gianmaria, Tiziana Cantile, Brunella Alcidi, Marco Coda, Aniello Ingenito, Armando Zarrelli, Giovanni Di Fabio, and Antonino Pollio. "Is Stevia rebaudiana Bertoni a Non Cariogenic Sweetener? A Review", *Molecules*, 2015.

Publication

<1 %

33

Alagarsamy Karthikeyan, Ramakrishnan Rameshkumar, Nallusamy Sivakumar, Issa Al Amri et al. "Antibiofilm Activity of *Dendrophthoe falcata* against Different Bacterial Pathogens", *Planta Medica*, 2012

Publication

<1 %

34 Hajera Khanam, Nishat Fatima. <1 %
"Pharmacological Screening for Burn Wound
Healing Potential of Traditional PlantCassia
fistuliain Rats", JOURNAL OF DRUG VIGILANCE
AND ALTERNATIVE THERAPIES, 2021
Publication

35 Hendry Rusdy, Diah HI Damanik. <1 %
"Antibacterial activity of Betadine (Jatropha
multifida L.) stem extract on Pseudomonas
aeruginosa growth in vitro", F1000Research,
2022
Publication

36 Indriyani Nur, Santi ., Agus Kurnia. <1 %
"Efficacy of
Guava (Psidium guajava) Leaves Extract to
Prevent Vibriosis in White Shrimp
(Litopenaeus vannamei)", Research Journal of
Medicinal Plants, 2019
Publication

37 Mendoza Monroy Ulises. <1 %
"Efecto sobre el
crecimiento de la bacteria cariogénica
Streptococcus mutans del extracto
metanólico de la corteza de Bursera simaruba
: actividad anticariogénica del lignano
picropoligamaina", TESIUNAM, 2011
Publication

38 Mónica Lilian Pérez-Ochoa, José Luis Chávez-
Servia, Araceli Minerva Vera-Guzmán, Elia
Nora Aquino-Bolaños et al. "Chapter 5
<1 %

Medicinal Plants Used by Indigenous Communities of Oaxaca, Mexico, to Treat Gastrointestinal Disorders", IntechOpen, 2019

Publication

39

Nining Kurniati. "Inhibitory Power of Papaya Leaves to the Adhesion of Streptococcus Mutans Bacteria to Neutrophils", Biomedical and Pharmacology Journal, 2021

Publication

40

Patryk Oleszczuk, Stanislaw Baran. "KINETICS OF PAHS LOSSES AND RELATIONSHIPS BETWEEN PAHS PROPERTIES AND PROPERTIES OF SOIL IN SEWAGE SLUDGE-AMENDED SOIL", Polycyclic Aromatic Compounds, 2007

Publication

41

Shabnam Mahernia, Malihe Hassanzadeh, Niusha Sharifi, Bitra Mehravi, Fariba Paytam, Mehdi Adib, Massoud Amanlou. "Structure-based pharmacophore design and virtual screening for novel potential inhibitors of epidermal growth factor receptor as an approach to breast cancer chemotherapy", Molecular Diversity, 2017

Publication

42

Yaschilal Muche Belayneh, Gedefaw Getnet Amare, Birhanu Geta Meharie, Zemene Demelash Kifle. "Evaluation of the

<1 %

<1 %

<1 %

<1 %

antiulcerogenic activity of hydromethanol extracts of *Solanum incanum* L. (Solanaceae) leaves and roots in mice; single and repeated dose study", *Metabolism Open*, 2021

Publication

43

Abu Bakar Siddique, Syed Mohammad Mizanur Rahman, Mohammad Amzad Hossain, Mohammad Amzad Hossain, Mohammad Abdur Rashid. "Phytochemical screening and comparative antimicrobial potential of different extracts of *Stevia rebaudiana* Bertoni leaves", *Asian Pacific Journal of Tropical Disease*, 2014

<1 %

Publication

44

Ahmad Gholami, Fariborz Moazami, Vahid Mehrabi, Yasamin Ghahramani. "Evaluation of the Antibacterial and Antifungal Effects of ProRoot MTA and Nano-fast Cement: An In Vitro Study", *The Journal of Contemporary Dental Practice*, 2020

<1 %

Publication

45

Arif Azhari, Muhammad Yogi Saputra, Muhammad Raffi Zakaria, Debora Silalahi et al. "Potential Antioxidant Constituent from Leaf of *Rhizophora apiculata* an Typical Mangrove at Lempasing, South Lampung Coast", *Stannum : Jurnal Sains dan Terapan Kimia*, 2022

<1 %

Publication

46

Benjamin Bhunu, Ruvimbo Mautsa, Stanley Mukanganyama. "Inhibition of biofilm formation in *Mycobacterium smegmatis* by *Parinari curatellifolia* leaf extracts", BMC Complementary and Alternative Medicine, 2017

Publication

<1 %

47

Mohammed Abdalbasit A. Gasmalla, Ruijin Yang, Xiao Hua. "Extraction of rebaudioside-A by sonication from *Stevia rebaudiana* Bertoni leaf and decolorization of the extract by polymers", Journal of Food Science and Technology, 2015

Publication

<1 %

48

Sedghi, M, and S Gholi-Toluie. "Influence of Salicylic Acid on the Antimicrobial Potential of *Stevia (Stevia rebaudiana* Bertoni, Asteraceae) Leaf Extracts against Soybean Seed-Borne Pathogens", Tropical Journal of Pharmaceutical Research, 2014.

Publication

<1 %

49

Siswo, Yun, Abdiyani. "Assessing Vegetation Composition and the Indicator Species Around Water Source Areas in a Pine Forest Plantation: A Case Study from Watujali and Silengkong Catchments, Kebumen, Indonesia", Forests, 2019

Publication

<1 %

50

Smitha Chenicheri, Usha R, Rajesh Ramachandran, Vinoy Thomas, Andrew Wood. "Insight into Oral Biofilm: Primary, Secondary and Residual Caries and Phyto-Challenged Solutions", The Open Dentistry Journal, 2017

Publication

<1 %

51

Vijayakumar Sundaram, Selvaraju Sadhasivam, Sivaraj Chandrasekaran, Raaman Nanjian, Arjun Pandian. "Strobilanthes heyneanus root extract as a potential source for antioxidant and antimicrobial activity", Future Journal of Pharmaceutical Sciences, 2021

Publication

<1 %

52

Audrey-Ann Dumont, Lauralyne Dumont, Delong Zhou, Hugo Giguère et al. "Cardiomyocyte-specific Srsf3 deletion reveals a mitochondrial regulatory role", Cold Spring Harbor Laboratory, 2020

Publication

<1 %

53

S. KHURSHID, A. JAVAID, A. SHOAIB, S. JAVED, U. QAISAR. "Antifungal Activity of Aerial Parts of Cenchrus pennisetiformis Against Fusarium oxysporum f. sp. lycopersici", Planta Daninha, 2018

Publication

<1 %

54

A. B. Falowo, V. Muchenje, A. Hugo, O. A. Aiyegoro, P. O. Fayemi. " Antioxidant activities

<1 %

of L. and L. leaf extracts and their effects on oxidative stability of ground raw beef during refrigeration storage ", CyTA - Journal of Food, 2016

Publication

55

Shilpa Sharma, Swati Gupta, Deepa Kumari, Shanker Lal Kothari, Rohit Jain, Sumita Kachhwaha. "Exploring Plant Tissue Culture and Steviol Glycosides Production in Stevia rebaudiana (Bert.) Bertoni: A Review", Agriculture, 2023

Publication

<1 %

56

Sudesh Kumar Yadav, Praveen Guleria. "Steviol Glycosides from Stevia: Biosynthesis Pathway Review and their Application in Foods and Medicine", Critical Reviews in Food Science and Nutrition, 2012

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off