

The Effect of Pegagan (*Centella Asiatica* (L.) Urban) Ethanol Extracts on Hippocampal PSD-95 Protein Expression in Male Wistar Rats

Adibah Ferhad¹, Auliyani Andam Suri¹, Astri Handayani¹, Sri Redjeki^{2*}, Ria Kodariah³

¹Biomedical Science Graduate Study Program, Faculty of Medicine, Universitas Indonesia

²Department of Physiology, Faculty of Medicine, Universitas Indonesia

³Department of Anatomic Pathology, Faculty of Medicine, Universitas Indonesia

ABSTRACT

One effort to overcome the decline in memory function is through herbal medicine. Pegagan (*Centella asiatica* (L.) Urban) contain the active components of triterpenoid and flavonoids, has been known to be able to improve memory function. Synaptic plasticity is the basis of memory formation which is strongly influenced by synapse proteins such as PSD-95. Loss of PSD-95 protein can cause memory function decline. This study aims to determine the effect of 70% ethanol extract of pegagan toward PSD-95 protein expression on hippocampus of male Wistar rat. Eighteen male Wistar rats were randomly divided into 3 groups, 6 rats/group: group (1) given ethanol extract of pegagan with dose 300 mg / kgBW / day (CA300), (2) given ethanol extract of pegagan with dose 600 mg / kgBW / day (CA600), and (3) control group (K), given daily aquadest. All three groups were treated for 28 consecutive days. At the end of the treatment period, rats were decapitated and the hippocampus was isolated from the brain. Analysis of protein was done by immunohistochemical method. Statistical analysis was performed by One Way ANOVA parametric test followed by Post-Hoc Bonferroni. The results showed that there was no significant difference between group K and group CA300 ($P = 0.123$), whereas there were significant differences between CA600 group and K group and CA300 group ($P = 0.000$). From this research, it can be concluded that 70% ethanol extract of pegagan with dose 600mg / kgBW / day can increase expression of PSD-95 protein on hippocampus of male Wistar rat.

Key words: *Post Synaptic Density-95*; memory function; flavonoids; *Centella asiatica* (L.) Urban.

INTRODUCTION

Aging is a natural biological process characterized by a decrease in physiological functions of the body including a decrease in memory function in the brain. (Shivarama Shetty and Sajikumar, 2017, Bettio *et al.*, 2017) Decreasing memory function can occur in the normal aging process and in the aging process accompanied by degenerative diseases. (Bettio *et al.*, 2017) Memory is the ability to record the life experiences so that affects a person's attitude or habit according to environmental conditions (Jared, 2010). The process of memory formation occurs through cellular mechanisms, namely the change in synapse strength called synapse plasticity (Lu *et al.*, 2008, Leal *et al.*, 2015). Plasticity at the excitatory synapse can be mediated at the level of pre-synapse and post-synapse (Leal *et al.*, 2015). Postsynaptic density-95 (PSD-95) is the main scaffold protein in the post-synapse density at the excitatory synaptic glutamate. PSD-95 is a member of the membrane-associated guanylate kinase (MAGUK) (Zhang *et al.*, 2013). PSD-95 plays a role in maintaining

and modulating synapse strength (Chen *et al.*, 2015). PSD-95 is often used as a marker of synapse at the post-synapse and is closely related to memory function (Nelson and Alkon, 2015). PSD-95 plays a role in regulating post-synaptic receptors such as NMDA receptors and AMPA receptors. (Han and Kim, 2008) The long-term potentiation mechanism related to memory formation in the hippocampus is influenced by the delivery of AMPA receptors to the post-synapse membrane and this PSD-95 can control the number of AMPA receptors in the post-synapse membrane. (Ehrlich and Malinow, 2004) The loss of PSD-95 protein that occurs along with increasing age is closely related to a decrease in memory function (Ojo *et al.*, 2012).

One of the effort to overcome the decrease in memory function is through herbal medication. Pegagan plant (*Centella asiatica* (L.) Urban) is a tropical medicinal plant from *Apiaceae* family originating from Southeast Asia including Indonesia. This pegagan leaf is edible, yellowish green, thin, with long petioles, and has a distinctive shape which is kidney-shaped, jagged, rounded, or oval-elliptical with seven leaf bones. These plants grow horizontally with reddish green stolon that

*Corresponding author : Sri Redjeki
Email : djeki.pras@gmail.com

merge each other and rooted underground (Orhan, 2012). *Pegagan* or *antan* is often found in open places, on moist and fertile soil such as in rice fields, in grasslands, on the edge of trenches and on the roadside (Ramadhan *et al.*, 2015). There are two main components in *pegagan*, namely triterpenoids and flavonoids. Both components function as antioxidants and are neuroprotective (Sari and Rochmah, 2015, Orhan, 2012, Vauzour *et al.*, 2008). *Pegagan* has long been used in Ayurveda medicine and traditional Chinese medicine to improve memory function (Lokanathan *et al.*, 2016). Previous research has shown that the distribution of ethanol extract of *Centella asiatica* for 8 weeks at a dose of 300 mg/kg.BW increased the cognitive function of male Wistar rats aged 2 months.(Mirza I, 2013) It is not known whether this increase in cognitive function is related to proteins that underlie the memory formation.

The decrease memory function in humans occurs early in adulthood, that is, since the age of 20 (Marlatt *et al.*, 2012). Therefore, this study was conducted to determine the effect of *pegagan* ethanol extract in increasing the expression of PSD-95 protein as a form of prevention of decrease memory function in hippocampus of Wistar male rats aged 6 months, where that age when converted to humans is equal to 18 (Andreollo *et al.*, 2012).

METHODOLOGY

Materials

Test drug materials

Simplicia of *pegagan* plant (*Centella asiatica* (L.) Urban) obtained from Pusat Studi Biofarmaka Institut Pertanian Bogor (IPB). Extraction of the materials was conducted at Balai Penelitian Tanaman Rempah dan Obat (Balittro) Bogor. *Pegagan* is extracted with 70% ethanol using maceration method. After the materials are extracted, a qualitative test is conducted with the KLT method and quantitative test is conducted using the spectrophotometric method and TLC Scanner to find out the active materials of the *pegagan* ethanol extract. The design and method of this research have passed the ethical review by Komite Etik Penelitian Kesehatan Fakultas Kedokteran Universitas Indonesia with an ethical number: 824/UN2.FI/ETIK/2016.

The subject of research

The subjects of this study were experimental animals Wistar strain male rats (*Rattus norvegicus*), 6 months old with an initial weight of 300 - 400 grams, healthy and active. The research subjects were obtained from Biofarma-Bandung. The treatment of experimental animals

was conducted at the Laboratorium Departemen Patologi Anatomi FKUI. Before and during the treatment, animal health was maintained so as not to get sick. Experimental animals are given standard food and drinks in *ad libitum*. The experimental animals is maintained in husk-bottomed cages that are equipped with feed containers and drinking water bottles. The animal cages are kept clean and arranged 12 hours bright and 12 hours dark. The ambient temperature is maintained at 25 + 1°C. Other things in the experiment were adjusted to the ethical code of commission for handling and using of experimental animals. The sample of this study was hippocampus tissue of male Wistar rats which was the subject of the research.

Materials and equipment for checking proteins

The procedure for examining the PSD-95 protein is conducted at Laboratorium Imunologi Departemen Patologi Anatomi FKUI using immunohistochemical methods (IHK). Materials and equipment used include xylene (Merck), ethanol (Merck) (with absolute concentration, 96%, 80% and 70%), Phosphate Buffer Saline (PBS) (Biomatix) with a pH of 7.4, sodium citrate buffer pH 6.0, Hydrogen peroxide (H2O2) 35%, primary antibody anti-PSD95 (Abcam), secondary antibodies Universal HRP Polymer (Neopoly), detection kit (Biogear), Hematoxylin Meyer, saturated Lithium carbonate (5%), entelan, super PAP pen (Biogear), and deionized water. While the equipment used include analytical scales, microtome, Aurona microscope slides (positively charged), cover glass, water bath, drying oven, slide warmer, Biocare's Decloaking Chamber, pH meter (Schoot), microscope (Olympus BX51) with a digital camera (Olympus).

Method

Preparation of test drug materials

The dried *pegagan* simplicia was extracted by maceration method at Balittro Bogor. 5 kg of dried *pegagan* simplicia is mashed using a grinder until it becomes powder. *Pegagan* powder was then weighed and obtained 4 kg of powder and then put into a stainless container and put in the solvent, 70% ethanol mixed with mixer for 3 hours. Subsequently precipitated over 24 hours. Thereafter filtered using filter paper. The filter results are concentrated by rotary evaporator at a temperature of 50°C until evaporation is complete. 326.4 gr thick extract was obtained and the rendement was 8.16% with a moisture content of 17.39%.

After extraction, phytochemical tests were conducted using the KLT qualitative method and

the results were obtained that *pegagan* extract contained alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides. Then proceed with a quantitative test to determine the percentage of active content of *pegagan* extract. Quantitative tests use spectrophotometric methods to determine tannin and flavonoids levels, which amounted to 0.22% and 0.44%. Quantitative tests use TLC Scanner method to determine saponin levels and asiaticoside levels, which amounted to 1.25% and 0.65%.

The distribution of *pegagan* extract

Experimental animals are weighed to determine the amount of extract to be given according to the dosage range. The preparation is prepared to fulfill the needs of the distribution extract. Dilution of the extract preparation was prepared for 7 days of treatment. This is conducted to avoid damage if the preparation is stored more than 7 days. After dilution, the preparation is stored in a refrigerator at 4°C. After 7 days, preparations are made for the next day. Before being given to experimental animals, the preparation extract was left for a moment at room temperature, then stirred well so that it was mixed and there was no sediment to facilitate the rounding process in the experimental animals.

Experimental animals are divided randomly into three groups, namely: (1) the control group (K) were given aquadest; (2) the group given the extract of *pegagan* with the dose of 300 mg/kg. W/day (CA 300); (3) the group was given a *pegagan* extract with a dose of 600 mg / kg.BW/day (CA 600). Each treatment towards every group was given once a day for 28 consecutive days at the same time in the morning, given orally using a stomach tube.

Series of immunohistochemistry procedures and protein checks

At the end of the treatment the research subjects were decapitated. Then the hippocampal tissue was isolated from the brain and put into 10% formalin buffer for 24 hours. Each treatment group was taken for each of the 6 hippocampal tissue from 6 experimental animals. Next, paraffin blocks were made and coronal directional slices were made with a thickness of 4 micrometers and floated in a water bath. Then the pieces are placed on coated slides. Liquids on the slides are dried, then the slides are put into drying oven for 30 – 60 minutes with temperature 37°C (Spencer and Bancroft, 2008).

The CPI discharge procedure used in this study is the Neopoly IHC procedure which is the standard procedure for Biogear Polymer Neopoly

Detection Kit. From the optimization results obtained, optimal dilution for PSD-95 protein is a concentration of 1:500. After the CPI procedure is completed, the expression percentage measurement of PSD-95 protein is done by observing slides under a binocular light microscope (Olympus BX51) with 400x magnification and the image is taken with a built-in camera (Olympus) on the microscope. For each slide five images are taken representing five visual fields (2560x1920 pixels) in the CA1 area of the hippocampal tissue. The resulting images are calculated using ImageJ program with the IHC Profiler plug-in to produce intensity values automatically for each image. Then, the results are calculated using IHC Optical Density Score method.

The optical density data of the PSD-95 protein that has been obtained is then carried out with statistical analysis using SPSS 17.0 computer program. Data were analyzed using one way ANOVA test then Post-Hoc Bonferroni test with 95% confidence level.

RESULTS AND DISCUSSION

One of the typical characteristics of aging is loss of memory function, especially in the hippocampus. This is related to changes in synapse morphology including loss of pre-synapse and post-synapse proteins, and progressive loss of synapse's density. (Ojo *et al.*, 2012) The hippocampus is located at the edge of the temporal lobe of the cerebral cortex consisting of Cornu Ammonis and dentate gyrus. Cornu ammonis (CA) consists of four parts namely CA1, CA2, CA3 and CA4. But two important Cornu Ammonis are CA1 and CA3. CA1 consists of pyramidal cells and plays an important role in matching information obtained from CA3. (Anand and Dhikav, 2012, Bear *et al.*, 2016) The development process of the human brain, including synapse density and synaptogenesis has started since inside the womb. Synaptogenesis is known to have occurred since the 34th week of pregnancy where nearly 40,000 new synapses are formed every second and continue until after birth, and begin to decline when entering adolescence period. (Tau and Peterson, 2010) Various studies also found that memory function began to decline after turning 20 years and goes on until 80 years old. (Marlatt *et al.*, 2012) Based on that theory, the age of experimental animals is set at 6 months which is equivalent to 18 years in humans. The age determination for experimental animals is in accordance with the research objectives to find *pegagan* extract can be used as an action to prevent decreased memory function.




Figure 1. Revitalized PSD-95 CPI in the CA1 area of hippocampal tissue in male Wistar rats. A. negative control; B. control group; C. CA300 group; D. CA600 group. 400X magnification. (Arrow shows the expression of PSD-95 protein in hippocampus pyramidal cells that are browned in the pyramidal cell cytoplasm). Experimental animals were decapitated on the 29th day and the hippocampal tissues were isolated, then paraffin blocks were made and then cut to 4 μ m. Recovering CPI is carried out with anti-PSD95 primary antibodies (Abcam), Universal HRP Polymer (Neopoly) secondary antibody.




Figure 2. Comparison of mean values of optical density of PSD-95 proteins between the three groups. *The results of statistical analysis using the One Way Anova test followed by the Bonferroni Post-Hoc test; there was a significant difference between the CA600 group and the control group ($p = 0.000$) and CA300 ($p = 0.000$). Significance * $p < 0.05$. The data represent 6 experimental animals per group. Data in mean values, error bars show standard deviation (Remarks: Kolmogorov-Smirnov test results shows data is not normally distributed, but the transformation is successful).

In this study, an examination of post synapse protein PSD-95 was examined in the hippocampal CA1 area of a male Wistar Rats. In Figure 1, the results of CPI output are shown on the sample network of control group, CA300, and CA600. Figure 1A shows a negative control where the expression of PSD-95 protein does not appear. Expression of PSD-95 protein in the CA1 region of male Wistar rats was seen to be the weakest in the control group (Figure 1B) and the strongest expression of PSD-95 protein was seen in the CA600 group (Figure 1D).

The mean optical density scores comparison between PSD-95 proteins in the three groups is shown in Figure 2. The highest mean value was found in the CA600 group (2.46 ± 0.16), followed by the CA300 group (1.83 ± 0.19) and the lowest was the control group (1.64 ± 0.09). Statistical analysis then carried out, with One Way Anova parametric test followed by Bonferroni Post-Hoc test. Based on the results of statistical analysis, it was found that there were significant differences between the CA600 group and the control group ($p = 0.000$) and CA300 group ($p = 0.000$), while the CA300 group and the control group does not differ significantly ($p = 0.123$).

The result of this study stated that administration of 70% *pegagan* ethanol extract can increase the expression of PSD-95 protein in the hippocampal CA1 area of male Wistar rats. The results of this study are supported by research which is conducted by Gray *et al.*, that administration of *pegagan* extract in mice can increase the synaptic gene expression of PSD-95 in the hippocampus (Gray *et al.*, 2016). Decision for giving 70% ethanol extract is because 70% ethanol is best in dissolving asiaticosides contained in *pegagan*. (Salamah and Farahana, 2014)

Pegagan plant has many benefits including anti-inflammatory, antimicrobial, antifungal, antidepressant, antioxidant and anticancer properties (Lokanathan *et al.*, 2016) as well as improving memory function and is neuroprotective (Orhan, 2012). The two main components in *pegagan* are triterpenoids and flavonoids. Triterpenoids consists of active components, namely asiatic acid and asiaticoside (Sari and Rochmah, 2015).

In this study it was stated that distributing the *Pegagan* ethanol extract containing asiaticoside for 28 consecutive days could increase the expression of PSD-95 in the male hippocampus of Wistar rats. The results of this study are supported by previous studies which stated that administration of asiaticoside for four consecutive weeks can increase the expression of PSD-95 in the hippocampus (Luo *et al.*, 2015).

CONCLUSION

The effect of giving *pegagan* ethanol extract (dose 600 mg / kg.BW) was better in increasing PSD-95 protein expression in the hippocampus of male Wistar rats compared to 300 mg / kg.BW dose and the control group.

ACKNOWLEDGEMENTS

This research is supported through 2017 Thesis and Dissertation Scholarship by Lembaga Pengelola Dana Pendidikan (LPDP), Ministry of Finance of the Republic of Indonesia.

REFERENCES

- Anand, K. S. & Dhikav, V. 2012. Hippocampus in health and disease: An overview. *Annals of Indian Academy of Neurology*, 15, 239.
- Andreollo, N. A., Santos, E. F. d., Araújo, M. R. & Lopes, L. R. 2012. Rat's age versus human's age: what is the relationship? *ABCD. Arquivos Brasileiros de Cirurgia Digestiva (São Paulo)*, 25, 49-51.
- Bear, M. F., Connors, B. W. & Paradiso, M. A. 2016. *Neuroscience: Exploring the Brain*. , Philadelphia, Wolters Kluwer.
- Bettio, L. E. B., Rajendran, L. & Gil-Mohapel, J. 2017. The effects of aging in the hippocampus and cognitive decline. *Neurosci Biobehav Rev*, 79, 66-86.
- Chen, X., Levy, J. M., Hou, A., Winters, C., Azzam, R., Sousa, A. A., Leapman, R. D., Nicoll, R. A. & Reese, T. S. 2015. PSD-95 family MAGUKs are essential for anchoring AMPA and NMDA receptor complexes at the postsynaptic density. *Proc Natl Acad Sci U S A*, 112, E6983-92.
- Ehrlich, I. & Malinow, R. 2004. Postsynaptic density 95 controls AMPA receptor incorporation during long-term potentiation and experience-driven synaptic plasticity. *J Neurosci*, 24, 916-27.
- Gray, N. E., Harris, C. J., Quinn, J. F. & Soumyanath, A. 2016. Centella asiatica modulates antioxidant and mitochondrial pathways and improves cognitive function in mice. *J Ethnopharmacol*, 180, 78-86.
- Han, K. & Kim, E. 2008. Synaptic adhesion molecules and PSD-95. *Prog Neurobiol*, 84, 263-83.
- Jared, S. R. 2010. Enhancement of memory in rats with Centella asiatica. *Biomedical Research*, 21.
- Leal, G., Afonso, P. M., Salazar, I. L. & Duarte, C. B. 2015. Regulation of hippocampal synaptic plasticity by BDNF. *Brain Res*, 1621, 82-101.
- Lokanathan, Y., Omar, N., Puzi, N. N. A., Saim, A. & Idrus, R. H. 2016. Recent updates in

- neuroprotective and neuroregenerative potential of *Centella asiatica*. *The Malaysian journal of medical sciences: MJMS*, 23, 4.
- Lu, Y., Christian, K. & Lu, B. 2008. BDNF: a key regulator for protein synthesis-dependent LTP and long-term memory? *Neurobiol Learn Mem*, 89, 312-23.
- Luo, L., Liu, X. L., Mu, R. H., Wu, Y. J., Liu, B. B., Geng, D., Liu, Q. & Yi, L. T. 2015. Hippocampal BDNF signaling restored with chronic asiaticoside treatment in depression-like mice. *Brain Res Bull*, 114, 62-9.
- Marlatt, M. W., Potter, M. C., Lucassen, P. J. & van Praag, H. 2012. Running throughout middle-age improves memory function, hippocampal neurogenesis, and BDNF levels in female C57BL/6J mice. *Dev Neurobiol*, 72, 943-52.
- Mirza I, R. H., Khomsan A, Marliyati SA, Damayanthi E, Winarto A. 2013. Pengaruh ekstrak etanol daun pegagan (*Centella asiatica* (L.) urban) terhadap gambaran darah, aktivitas, dan fungsi kognitif tikus. *Jurnal Kedokteran Hewan*, 7.
- Nelson, T. J. & Alkon, D. L. 2015. Molecular regulation of synaptogenesis during associative learning and memory. *Brain Res*, 1621, 239-51.
- Ojo, B., Rezaie, P., Gabbott, P. L., Davies, H., Colyer, F., Cowley, T. R., Lynch, M. & Stewart, M. G. 2012. Age-related changes in the hippocampus (loss of synaptophysin and glial-synaptic interaction) are modified by systemic treatment with an NCAM-derived peptide, FGL. *Brain Behav Immun*, 26, 778-88.
- Orhan, I. E. 2012. *Centella asiatica* (L.) Urban: From Traditional Medicine to Modern Medicine with Neuroprotective Potential. *Evid Based Complement Alternat Med*, 2012, 946259.
- Ramadhan, N. S., Rasyid, R. & Syamsir, E. 2015. Daya Hambat Ekstrak Daun Pegagan (*Centella asiatica*) yang Diambil di Batusangkar terhadap Pertumbuhan Kuman *Vibrio cholerae* secara In Vitro. *Jurnal Kesehatan Andalas*, 4.
- Salamah, N. & Farahana, L. 2014. Uji Aktivitas Antioksidan Ekstrak Etanol Herba Pegagan (*Centella asiatica* (L.) Urb) dengan Metode Fosfomolibdat. *Pharmaciana*, 4.
- Sari, D. C. R. & Rochmah, M. A. 2015. The effects of ethanol extracts of *Centella asiatica* leaf on serial serum brain derived neurotrophin factor (bdnf) concentration of rats (sprague dawley) following chronic stress. *KnE Life Sciences*, 2, 159.
- Shivarama Shetty, M. & Sajikumar, S. 2017. 'Tagging' along memories in aging: Synaptic tagging and capture mechanisms in the aged hippocampus. *Ageing Res Rev*, 35, 22-35.
- Spencer, L. T. & Bancroft, J. D. 2008. Microtomy: paraffin and frozen. Dalam: Bancroft JD, Gamble M. *Theory and practice of histological techniques*. 6th ed. Philadelphia: Elsevier.
- Tau, G. Z. & Peterson, B. S. 2010. Normal development of brain circuits. *Neuropsychopharmacology*, 35, 147-68.
- Vauzour, D., Vafeiadou, K., Rodriguez-Mateos, A., Rendeiro, C. & Spencer, J. P. 2008. The neuroprotective potential of flavonoids: a multiplicity of effects. *Genes Nutr*, 3, 115-26.
- Zhang, J., Lewis, S. M., Kuhlman, B. & Lee, A. L. 2013. Supertertiary structure of the MAGUK core from PSD-95. *Structure*, 21, 402-13.

LEMBAR
AWAT SEBIDAN
Jurnal Nasional

Judul Jurnal Ilmiah (Artikel)	:	The Effect of Pegagan (<i>Centella Asiatica</i> (L.) Urban) Ethanol Extracts on Hippocampal PSD-95 Protein Expression in Male Wistar Rats
Penulis Jurnal Ilmiah	:	Adibah Ferhad
Identitas Jurnal Ilmiah	:	
a) Nama Jurnal	:	Trad. Med. J.
b) Nomor/Volume	:	Vol. 23 No.3 Tahun 2018
c) Edisi (Bulan/Tahun)	:	September – Desember 2018
d) Jumlah halaman	:	6 hal
e) ISSN/ISBN	:	1410-5918 ISSN-e: 2406-9086
f) Penerbit	:	Majalah Obat Tradisional (Trad.Med.J) Faculty of Pharmacy - Universitas Gadjah Mada
g) Url Asli	:	https://jurnal.ugm.ac.id/TradMedJ/article/view/33476/23403
h) Url Repository	:	
i) Url Review	:	

Kategori Publikasi Jurnal Ilmiah
(beri ✓ pada kategori yang tepat)

	Jurnal Ilmiah Internasional
x	Jurnal Ilmiah Nasional Terakreditasi
	Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi:

NO.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	11/02
2.	Linearitas	✓

II. Hasil Penilaian Peer Review:

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah					Nilai Akhir Yang diperoleh
	Inter nasional Bereputas	Inter nasional	Nasional Terakreditasi Sinta 2.	Nasional Tidak Terakreditasi	Nasional Terindeks DOAJ	
Kelengkapan dan kesesuaian isi Jurnal (10%)			10.			
Ruang Lingkup dan Kedalaman Pembahasan (30%)			28.			
Kecukupan dan Kemutakhiran Data/ Informasi dan Metodologi (30%)			30.			
Kelengkapan Unsur dan Kualitas Penerbit (30%)			30.			
Total=100%			98%			
Kontribusi Pengusul :	$60\% \times 24,5 = 14,7$					

Kontribusi Pengusul :

Komentar Peer Review :

1. Tentang Kelengkapan dan Kesesuaian Unsur *sesuai*
2. Tema baru dalam memahami *sesuai*
3. Kecukupan dan Kemutakhiran data serta Metodologi *sesuai*
4. Kelengkapan Unsur Kualitas Penerbit *sesuai*
5. Indikasi Plagiasi *11 %*
6. Kesesuaian Bidang Ilmu *sesuai*

Jakarta, 1 Oktober 2021
Penilai I



Nama : Dr. Tjebbo D. dy. Sp. Ort.
NIP/NIDN : 0325078005
Unit Kerja : WABEK I
Bidang Ilmu : Ortodonti
Jabatan/Pangkat : Rektor

LEMBAR
AWAT SEBIDAN
Jurnal Nasional

Judul Jurnal Ilmiah (Artikel)	:	The Effect of Pegagan (<i>Centella Asiatica</i> (L.) Urban) Ethanol Extracts on Hippocampal PSD-95 Protein Expression in Male Wistar Rats
Penulis Jurnal Ilmiah	:	Adibah Ferhad
Identitas Jurnal Ilmiah	:	
a) Nama Jurnal	:	Trad. Med. J.
b) Nomor/Volume	:	Vol. 23 No.3 Tahun 2018
c) Edisi (Bulan/Tahun)	:	September – Desember 2018
d) Jumlah halaman	:	6 hal
e) ISSN/ISBN	:	1410-5918 ISSN-e: 2406-9086
f) Penerbit	:	Majalah Obat Tradisional (Trad.Med.J) Faculty of Pharmacy - Universitas Gadjah Mada
g) Url Asli	:	https://jurnal.ugm.ac.id/TradMedJ/article/view/33476/23403
h) Url Repository	:	
i) Url Review	:	

Kategori Publikasi Jurnal Ilmiah
(beri ✓ pada kategori yang tepat)

	Jurnal Ilmiah Internasional
X	Jurnal Ilmiah Nasional Terakreditasi
	Jurnal Ilmiah Nasional Tidak Terakreditasi

1. Hasil Penilaian Validasi:

NO.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	
2.	Linearitas	Linear

II. Hasil Penilaian Peer Review:

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah					Nilai Akhir Yang diperoleh
	Inter nasional Bereputas	Inter nasional	Nasional Terakreditasi max 20	Nasional Tidak Terakreditasi	Nasional Terindeks DOAJ	
Kelengkapan dan kesesuaian isi Jurnal (10%)			9,5%			1,9
Ruang Lingkup dan Kedalaman Pembahasan (30%)			29%			5,8
Kecukupan dan Kemutakhiran Data/ Informasi dan Metodologi (30%)			29%			5,8
Kelengkapan Unsur dan Kualitas Penerbit (30%)			28%			5,6
Total=100%						19,1
Kontribusi Pengusul :	Penulis Pertama					

Komentar Peer Review :

1. Tentang Kelengkapan dan Kesesuaian Unsur Sesuai dengan panduan jurnal
2. Tema baru dalam memahami Tema baru dalam memahami efek dari tanaman pegagan yang diekstraksi terhadap eksresi Hippocampal PSD-95 dikuti
3. Kecukupan dan Kemutakhiran data serta Metodologi referensi relatif baru dan metodologi dapat ditentang jawabkan
4. Kelengkapan Unsur Kualitas Penerbit Tradisional Medicine jurnal, jurnal nasional terakreditasi
5. Indikasi Plagiasi tidak ada plagiasi
6. Kesesuaian Bidang Ilmu Sesuai dengan ilmu perilaku

Jakarta,
Penilai II

21 September 2021
Dr. drg. Sari Dewiyanti SpKG

Nama : Dr. drg. Sari Dewiyanti SpKG
NIP/NIDN :
Unit Kerja : FKG UPPM (LB)
Bidang Ilmu : Konservasi gigi
Jabatan/Pangkat : Lektor Kepala

The Effect of Pegagan (*Centella Asiatica* (L.) Urban) Ethanol Extracts on Hippocampal PSD-95 Protein Expression in Male Wistar Rats

Adibah Ferhad¹, Iuliyani Andam Suri¹, Astri Handayani¹, Sri Redjeki^{2*}, Ria Kodariah³

¹Biomedical Science Graduate Study Program, Faculty of Medicine, Universitas Indonesia

²Department of Physiology, Faculty of Medicine, Universitas Indonesia

³Department of Anatomic Pathology, Faculty of Medicine, Universitas Indonesia

ABSTRACT

One effort to overcome the decline in memory function is through herbal medicine. Pegagan (*Centella asiatica* (L.) Urban) contain the active components of terpenoid and flavonoids, has been known to be able to improve memory function. Synaptic plasticity is the basis of memory formation which is strongly influenced by synapse proteins such as PSD-95. Loss of PSD-95 protein can cause memory function decline. This study aims to determine the effect of 70% ethanol extract of pegagan toward PSD-95 protein expression on hippocampus of male Wistar rat. Eighteen male Wistar rats were randomly divided into 3 groups, 6 rats/group: group (1) given ethanol extract of pegagan with dose 300 mg / kgBW / day (CA300), (2) given ethanol extract of pegagan with dose 600 mg / kgBW / day (CA600), and (3) control group (K), given distilled aquadest. All three groups were treated for 28 consecutive days. At the end of the treatment period, rats were decapitated and the hippocampus was isolated from the brain. Analysis of protein was done by immunohistochemical method. Statistical analysis was performed by One Way ANOVA parametric test followed by Post-Hoc Bonferroni. The results showed that there was no significant difference between group K and group CA300 ($P = 0.123$), whereas there were significant differences between CA600 group and K group and CA300 group ($P = 0.000$). From this research, it can be concluded that 70% ethanol extract of pegagan with dose 600mg / kgBW / day can increase expression of PSD-95 protein on hippocampus of male Wistar rat.

Key words: Post Synaptic Density-95; memory function; flavonoids; *Centella asiatica* (L.) Urban.

INTRODUCTION

Aging is a natural biological process characterized by a decrease in physiological functions of the body including a decrease in memory function in the brain. (Shivarama Shetty and Sajikumar, 2017, Bettio *et al.*, 2017) Decreasing memory function can occur in the normal aging process and in the aging process accompanied by degenerative diseases. (Bettio *et al.*, 2017) Memory is the ability to record the life experiences so that affects a person's attitude or habit according to environmental conditions (Jared, 2010). The process of memory formation occurs through cellular mechanisms, namely the change in synapse strength called synapse plasticity (Lu *et al.*, 2008, Leal *et al.*, 2015). Plasticity at the excitatory synapse can be mediated at the level of pre-synapse and post-synapse (Leal *et al.*, 2015). Postsynaptic density-95 (PSD-95) is the main scaffold protein in the post-synapse density at the excitatory synaptic glutamate. PSD-95 is a member of the membrane-associated guanylate kinase (MAGUK) (Zhang *et al.*, 2013). PSD-95 plays a role in maintaining

and modulating synapse strength (Chen *et al.*, 2015). PSD-95 is often used as a marker of synapse at the post-synapse and is closely related to memory function (Nelson and Alkon, 2015). PSD-95 plays a role in regulating post-synaptic receptors such as NMDA receptors and AMPA receptors. (Han and Kim, 2008) The long-term potentiation mechanism related to memory formation in the hippocampus is influenced by the delivery of AMPA receptors to the post-synapse membrane and this PSD-95 can control the number of AMPA receptors in the post-synapse membrane. (Ehrlich and Malinow, 2004) The loss of PSD-95 protein that occurs along with increasing age is closely related to a decrease in memory function (Ojo *et al.*, 2012).

One of the effort to overcome the decrease in memory function is through herbal medication. Pegagan plant (*Centella asiatica* (L.) Urban) is a tropical medicinal plant from *Apiaceae* family originating from Southeast Asia including Indonesia. This pegagan leaf is edible, yellowish green, thin, with long petioles, and has a distinctive shape which is kidney-shaped, jagged, rounded, or oval-elliptical with seven leaf bones. These plants grow horizontally with reddish green stolon that

*Corresponding author : Sri Redjeki
Email : djeki.pras@gmail.com

merge each other and rooted underground (Orhan, 2012). Pegagan or antanan is often found in open places, on moist and fertile soil such as in rice fields, in grasslands, on the edge of trenches and on the roadside (Ramadhan *et al.*, 2015). There are two main components in pegagan, namely triterpenoids and flavonoids. Both components function as antioxidants and are neuroprotective (Sari and Rochmah, 2015, Orhan, 2012, Vauzour *et al.*, 2008). Pegagan has long been used in Ayurveda medicine and traditional Chinese medicine to improve memory function (Lokanathan *et al.*, 2016). Previous research has shown that the distribution of ethanol extract of *Centella asiatica* for 8 weeks at a dose of 300 mg/kg.BW increased the cognitive function of male Wistar rats aged 2 months. (Mirza I, 2013) It is not known whether this increase in cognitive function is related to proteins that underlie the memory formation.

The decrease memory function in humans occurs early in adulthood, that is, since the age of 20 (Marlatt *et al.*, 2012). Therefore, this study was conducted to determine the effect of pegagan ethanol extract in increasing the expression of PSD-95 protein as a form of prevention of decrease memory function in hippocampus of Wistar male rats aged 6 months, where that age when converted to humans is equal to 18 (Andreollo *et al.*, 2012).

METHODOLOGY

Materials

Test drug materials

Simplicia of pegagan plant (*Centella asiatica* (L.) Urban) obtained from Pusat Studi Biofarmaka Institut Pertanian Bogor (IPB). Extraction of the materials was conducted at Balai Penelitian Tanaman Rempah dan Obat (Balittro) Bogor. Pegagan is extracted with 70% ethanol using maceration method. After the materials are extracted, a qualitative test is conducted with the KLT method and quantitative test is conducted using the spectrophotometric method and TLC Scanner to find out the active materials of the pegagan ethanol extract. The design and method of this research have passed the ethical review by Komite Etik Penelitian Kesehatan Fakultas Kedokteran Universitas Indonesia with an ethical number: 824/UN2.FI/ETIK/2016.

The subject of research

The subjects of this study were experimental animals Wistar strain male rats (*Rattus norvegicus*), 6 months old with an initial weight of 300 - 400 grams, healthy and active. The

research subjects were obtained from Biofarma-Bandung. The treatment of experimental animals

Sri Redjeki

was conducted at the Laboratorium Departemen Patologi Anatomi FKUI. Before and during the treatment, animal health was maintained so as not to get sick. Experimental animals are given standard food and drinks in *ad libitum*. The experimental animals is maintained in husk-bottomed cages that are equipped with feed containers and drinking water bottles. The animal cages are kept clean and arranged 12 hours bright and 12 hours dark. The ambient temperature is maintained at $25 \pm 1^\circ\text{C}$. Other things in the experiment were adjusted to the ethical code of commission for handling and using of experimental animals. The sample of this study was hippocampus tissue of male Wistar rats which was the subject of the research.

Materials and equipment for checking proteins

The procedure for examining the PSD-95 protein is conducted at Laboratorium Imunologi Departemen Patologi Anatomi FKUI using immunohistochemical methods (IHK). Materials and equipment used include xylene (Merck), ethanol (Merck) (with 15 solute concentration, 96%, 80% and 70%), Phosphate Buffer Saline (PBS) (Biomatix) with a pH of 7.4, sodium citrate buffer pH 6.0, Hydrogen peroxide (H2O2) 35%, primary antibody anti-PSD95 (Abcam), secondary antibodies Universal HRP Polymer (Neopoly), detection kit (Biogear), Hematoxilin Meyer, saturated Lithium carbonate (5%), entelan, super PAP pen (Biogear), and deionized water. While the equipment used include analytical scales, microtome, Aurona microscope slides (positively charged), cover glass, water bath, drying oven, slide warmer, Biocare's Decloaking Chamber, pH meter (Schoot), microscope (Olympus BX51) with a digital camera (Olympus).

Method

Preparation of test drug materials

The dried *pegagan* simplicia was extracted by maceration method at Balitro Bogor. 5 kg of dried *pegagan* simplicia is mashed using a grinder until it becomes powder. *Pegagan* powder was then weighed and obtained 4 kg of powder and then put into a stainless container and put in the solvent, 70% ethanol mixed with mixer for 3 hours. Subsequently precipitated over 24 hours. Thereafter filtered using filter paper. The filter results are concentrated by rotary evaporator at a temperature of 50°C until evaporation is complete. 326.4 gr thick extract was obtained and the rendement was 8.16% with a moisture content of 17.39%.

After extraction, phytochemical tests were conducted using the KLT qualitative method and

the result⁹ were obtained that *pegagan* extract contained alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides. Then proceed with a quantitative test to determine the percentage of active content of *pegagan* extract. Quantitative tests use spectrophotometric methods to determine tannin and flavonoids levels, which amounted to 0.22% and 0.44%. Quantitative tests use TLC Scanner method to determine saponin levels and asiaticoside levels, which amounted to 1.25% and 0.65%.

The distribution of *pegagan* extract¹⁴

Experimental animals are weighed to determine the amount of extract to be given according to the dosage range. The preparation is prepared to fulfill the needs of the distribution extract. Dilution of the extract preparation was prepared for 7 days of treatment. This is conducted to avoid damage if the preparation is stored more than 7 days. After dilution, the preparation is stored in a refrigerator at 4°C. After 7 days, preparations are made for the next day. Before being given to experimental animals, the preparation extract was left for a moment at room temperature, then stirred well so that it was mixed and there was no sediment to facilitate the rounding process in the experimental animals.

Experimental animals are divided randomly into three groups, namely: (1) the control group (K) were given aquadest; (2) the group given the extract of *pegagan* with the dose of 300 mg/kg. W/day (CA 300); (3) the group was given a *pegagan* extract with a dose of 600 mg / kg.BW/day (CA 600). Each treatment towards every group was given once a day for 28 consecutive days at the same time in the morning, given orally using a stomach tube.

Series of immunohistochemistry procedures and protein checks

At the end of the treatment the research subjects were decapitated. Then the hippocampal tissue was isolated from the brain and put into 10% formalin buffer for 24 hours. Each treatment group was taken for each of the 6 hippocampal tissue from 6 experimental animals. Next, paraffin blocks were made and coronal directional slices were made with a thickness of 4 micrometers and floated in a water bath. Then the pieces are placed on coated slides. Liquids on the slides are dried, then the slides are put into drying oven for 30 – 60 minutes with temperature 37°C (Spencer and Bancroft, 2008).

The CPI discharge procedure used in this study is the Neopoly IHC procedure which is the standard procedure for Biogear Polymer Neopoly

Sri Redjeki

Detection Kit. From the optimization results obtained, optimal dilution for PSD-95 protein is a concentration of 1:500. After the CPI procedure is completed, the expression percentage measurement of PSD-95 protein is done by observing slides under a binocular light microscope (Olympus BX51) with 400x magnification and the image is taken with a built-in camera (Olympus) on the microscope. For each slide five images are taken representing five visual fields (2560x1920 pixels) in the CA1 area of the hippocampal tissue. The resulting images are calculated using ImageJ program with the IHC Profiler plug-in to produce intensity values automatically for each image. Then, the results are calculated using IHC Optical Density Score method.

The optical density data of the PSD-95 protein that has been obtained is then carried out with statistical analysis using SPSS 17.0 Computer program. Data were analyzed using one way ANOVA test then Post-Hoc Bonferroni test with 95% confidence level.

RESULTS AND DISCUSSION

One of the typical characteristics of aging is loss of memory function, especially in the hippocampus. This is related to changes in synapse morphology including loss of pre-synapse and post-synapse proteins, and progressive loss of synapse's density. (Ojo *et al.*, 2012) The hippocampus is located at the edge of the temporal lobe of the cerebral cortex consisting of Cornu Ammonis and dentate gyrus. Cornu ammonis (CA) consists of four parts namely CA1, CA2, CA3 and CA4. But two important Cornu Ammonis are CA1 and CA3. CA1 consists of pyramidal cells and plays an important role in matching information obtained from CA3. (Anand and Dhikav, 2012, Bear *et al.*, 2016) The development process of the human brain, including synapse density and synaptogenesis has started since inside the womb. Synaptogenesis is known to have occurred since the 34th week of pregnancy where nearly 40,000 new synapses are formed every second and continue until after birth, and begin to decline when entering adolescence period. (Tau and Peterson, 2010) Various studies also found that memory function began to decline after turning 20 years and goes on until 80 years old. (Marlatt *et al.*, 2012) Based on that theory, the age of experimental animals is set at 6 months which is equivalent to 18 years in humans. The age determination for experimental animals is in accordance with the research objectives to find pegagan extract can be used as an action to prevent decreased memory function.




Figure 1. Revitalized PSD-95 CPI in the CA1 area of hippocampal tissue in male Wistar rats. A. negative control; B. control group; C. CA300 group; D. CA600 group. 400X magnification. (Arrow shows the expression of PSD-95 protein in hippocampus pyramidal cells that are browned in the pyramidal cell cytoplasm). Experimental animals were decapitated on the 29th day and the hippocampal tissues were isolated, then paraffin blocks were made and then cut to 4 μ m. Recovering CPI is carried out with anti-PSD95 primary antibodies (Abcam), Universal HRP Polymer (Neopoly) secondary antibody.




Figure 2. Comparison of mean values of optical density of PSD-95 proteins between the three groups. *The results of statistical analysis using the One Way Anova test followed by the Bonferroni Post-Hoc test; there was a significant difference between the CA600 group and the control group ($p = 0.000$) and CA300 ($p = 0.000$). Significance * $p < 0.05$. The data represent 6 experimental animals per group. Data in mean values, error bars show standard deviation (Remarks: Kolmogorov-Smirnov test results shows data is not normally distributed, but the transformation is successful).

In this study, an examination of post synapse protein PSD-95 was examined in the hippocampal CA1 area of a male Wistar Rats. In Figure 1, the results of CPI output are shown on the sample network of control group, CA300, and CA600. Figure 1A shows a negative control where the expression of PSD-95 protein does not appear. Expression of PSD-95 protein in the CA1 region of male Wistar rats was seen to be the weakest in the control group (Figure 1B) and the strongest expression of PSD-95 protein was seen in the CA600 group (Figure 1D).

The mean optical density scores comparison between PSD-95 proteins in the three groups is shown in Figure 2. The highest mean value was found in the CA600 group (2.46 ± 0.16), followed by the CA300 group (1.83 ± 0.19) and the lowest was the control group (1.64 ± 0.09). Statistical analysis then carried out, with One Way Anova parametric test followed by Bonferroni Post-Hoc test. Based on the results of statistical analysis, it was found that there were significant differences between the CA600 group and the control group ($p = 0.000$) and CA300 group ($p = 0.000$), while the CA300 group and the control group does not differ significantly ($p = 0.123$).

The result of this study stated that administration of 70% pegagan ethanol extract can increase the expression of PSD-95 protein in the hippocampal CA1 area of male Wistar rats. The results of this study are supported by research which is conducted by Gray *et al.*, that administration of pegagan extract in mice can increase the synaptic gene expression of PSD-95 in the hippocampus (Gray *et al.*, 2016). Decision for giving 70% ethanol extract is because 70% ethanol is best in dissolving asiaticosides contained in pegagan. (Salamah and Farahana, 2014)

Pegagan plant has many benefits including anti-inflammatory, antimicrobial, antifungal, antidepressant, antioxidant and anticancer properties (Lokanathan *et al.*, 2016) as well as improving memory function and is neuroprotective (Orhan, 2012). The two main components in pegagan are triterpenoids and flavonoids. Triterpenoids consists of active components, namely asiatic acid and asiaticoside (Sari and Rochmah, 2015).

In this study it was stated that distributing the Pegagan ethanol extract containing asiaticoside for 28 consecutive days could increase the expression of PSD-95 in the male hippocampus of Wistar rats. The results of this study are supported by previous studies which stated that administration of asiaticoside for four consecutive

weeks can increase the expression of PSD-95 in the hippocampus (Luo *et al.*, 2015).

Sri Redjeki

²
The Effect of Pegagan (*Centella Asiatica* (L.) Urban)

CONCLUSION

The effect of giving *pegagan* ethanol extract (dose 600 mg / kg.BW) was better in increasing PSD-95 protein expression in the hippocampus of male Wistar rats compared to 300 mg / kg.BW dose and the control group.

ACKNOWLEDGEMENTS

This research is supported through 2017 Thesis and Dissertation Scholarship by Lembaga Pengelola Dana Pendidikan (LPDP), Ministry of Finance of the Republic of Indonesia.

Jurnal Nasional

ORIGINALITY REPORT

11 %

SIMILARITY INDEX

%

INTERNET SOURCES

11 %

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

Astri Handayani, Sophie Yolanda, Ria Kodariah. "Centella asiatica ethanol extract increases hippocampal brain derived neurotrophic factor in male Wistar rats", *Universa Medicina*, 2018

Publication

2 %

2

N S Vinolina, R Sigalingging. "Phytochemical screening and biomass production of *Centella asiatica* (L.) Urb of Samosir – Indonesia accession cultivated on acid soil with different phosphorus treatments", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

1 %

3

Tsuyoshi Nishiguchi, Masaaki Iwata, Naofumi Kajitani, Akihiko Miura et al. "Stress increases blood beta - hydroxybutyrate levels and prefrontal cortex NLRP3 activity jointly in a rodent model", *Neuropsychopharmacology Reports*, 2021

Publication

1 %

4

Xiaohong Xu, Lingjing Ye, Qin Ruan.
"Environmental Enrichment Induces Synaptic
Structural Modification After Transient Focal
Cerebral Ischemia in Rats", Experimental
Biology and Medicine, 2009

Publication

1 %

5

Qian Li, Baojun Yang, Jianyu Zhu, Hao Jiang,
Jiaokun Li, Ruiyong Zhang, Wolfgang Sand.
"Comparative Analysis of Attachment to
Chalcopyrite of Three Mesophilic Iron and/or
Sulfur-Oxidizing Acidophiles", Minerals, 2018

Publication

1 %

6

Fuzhi Zhuang, Mei Li, Xin Gao, Yun Wang,
Dongdong Wang, Xing Ma, Tengfei Ma,
Shuling Gu. "The antidepressant-like effect of
alarin is related to TrkB-mTOR signaling and
synaptic plasticity", Behavioural Brain
Research, 2016

Publication

1 %

7

Manik Hapsara, Ahmed Imran, Timothy
Turner. "Chapter 3 E-Voting in Developing
Countries", Springer Science and Business
Media LLC, 2017

Publication

<1 %

8

Patiyus Agustiansyah, Rizal Sanif, Siti
Nurmaini, Irfannuddin, Legiran. "Epidemiology
and Risk Factors for Cervical Cancer",

<1 %

9

Y Koniyo, C Lumenta, A H Olli, R O S E Mantiri,
N Pasingi. " Nutrition of local wild edible fern
() leaves ", IOP Conference Series: Earth and
Environmental Science, 2021

Publication

<1 %

10

Adam Satria Rakatama, Andri Pramono, Retno
Yulianti. " The Antifungal Inhibitory
Concentration Effectiveness Test From
Ethanol Seed Arabica Coffee () Extract
Against The Growth Of Candida albicans
Patient Isolate With In Vitro Method ", Journal
of Physics: Conference Series, 2018

Publication

<1 %

11

Berta Henriquez, Fernando J. Bustos, Rodrigo
Aguilar, Alvaro Becerra, Felipe Simon, Martin
Montecino, Brigitte van Zundert. "Ezh1 and
Ezh2 differentially regulate PSD-95 gene
transcription in developing hippocampal
neurons", Molecular and Cellular
Neuroscience, 2013

Publication

<1 %

12

Deng, Jing, Juan Yang, Lan-Fang Wu, Kang-
Kang Yu, Dao-Ping Wang, Xiao-Sheng Yang,
and Fu-Xue Chen. "Structural Characterization
and Neurotrophic Activity Study of a

<1 %

Polysaccharide Isolated from *Campanumoea javanica*", Journal of Carbohydrate Chemistry, 2015.

Publication

13

Marcia Dewi Hartanto, Zita Arieselia, Rianto Setiabudy, Arini Setiawati, Ali Baziad. "Urinary 11-dehydro-thromboxane B2 and 2,3-dinor-6-keto-prostaglandin-F1 α in healthy post-menopausal and pre-menopausal women receiving aspirin 100 mg", Journal of Thrombosis and Thrombolysis, 2012

Publication

<1 %

14

Neng Tine Kartinah, Nisa Komara, Nuraini Diah Noviati, Syarifah Dewi et al. "Potential of Hibiscus sabdariffa linn. in managing Fgf21 resistance in diet-induced-obesity rats via miR-34a regulation", Research Square, 2020

Publication

<1 %

15

Dyah Kinasih Wuragil, Masdiana Chendrakasih Padaga, Aulanni'am Aulanni'am, Chanif Mahdi. "Study of goat milk yogurt for hypercholesterolemic rats (*Rattus norvegicus*) models", AIP Publishing, 2019

Publication

<1 %

16

Foqia Khalid, Muhammad Zaffar Hashmi, Nadia Jamil, Abdul Qadir. "Screening of Wetland Plants Subjected to Stress From

<1 %

Polychlorinated Biphenyls Contaminated Soils", Research Square Platform LLC, 2021

Publication

17

Juminten Saimin, Hendy Hendarto, Soetjipto Soetjipto. "The Effect of Tomato Juice in Increasing Ki-67 Expression and Epithelial Thickness on The Vaginal Wall of Menopausal Rats", The Indonesian Biomedical Journal, 2019

Publication

<1 %

18

Wadhwa, M., S. Sahu, P. Kumari, H. Kauser, K. Ray, and U. Panjwani. "Caffeine and modafinil given during 48h sleep deprivation modulate object recognition memory and synaptic proteins in the hippocampus of the rat", Behavioural Brain Research, 2015.

Publication

<1 %

19

Zhi-Wei Zhu. "Spatiotemporal expression of PSD-95 in Fmr1 knockout mice brain : Profile of PSD-95 in Fmr1 KO mice brain", Neuropathology, 06/2011

Publication

<1 %

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off